

ON A NIELSEN-THURSTON CLASSIFICATION THEORY ON CLUSTER MODULAR GROUPS

TSUKASA ISHIBASHI

ABSTRACT. We classify elements of a cluster modular group into three types. We characterize them in terms of fixed point property of the action on the tropical compactifications associated with the corresponding cluster ensemble. The characterization gives an analogue of the Nielsen-Thurston classification theory on the mapping class group of a surface.

CONTENTS

0. Introduction	1
1. Definition of the cluster modular groups	3
1.1. The cluster modular groups and the cluster ensembles	3
1.2. Cluster complexes	7
1.3. Tropical compactifications of positive spaces	8
2. Nielsen-Thurston types on cluster modular groups	9
2.1. Periodic classes	10
2.2. Cluster-reducible classes	12
2.3. Cluster-pA classes of special type: cluster Dehn twists	16
3. Basic examples: SL_2 -Teichmüller seeds	18
3.1. Definition of the seed	18
3.2. The SL_2 -Teichmüller seed on an marked hyperbolic surface is of Teichmüller type	21
3.3. Comparison with the Nielsen-Thurston classification of elements of the mapping class group	22
Appendix A. The seed of type L_k is of Teichmüller type	24
Appendix B. The positive real parts and the tropical spaces associated with the SL_2 -Teichmüller seed	25
B.1. Positive spaces and the Teichmüller spaces	26
B.2. Tropical spaces and the lamination spaces	28
References	30

0. INTRODUCTION

A *cluster modular group*, defined in [FG09], is a discrete group associated with a combinatorial data called a *seed*. An element of a cluster modular group is a finite sequence of permutations of vertices and *mutations*, which preserve the *exchange matrix* and induce a pair of non-trivial *cluster transformations*. It is known that, for each marked hyperbolic surface F , the cluster modular group associated with a certain seed includes the *mapping class group* of F as a subgroup of finite index. We call this seed the *SL_2 -Teichmüller seed* on F in this paper. Therefore it seems natural to ask whether a property known for mapping class groups holds for general cluster modular groups. In this paper we attempt to provide an analogue of the

Nielsen-Thurston theory [Thu88, FLP12] on mapping class groups, which classifies mapping classes into three types in terms of fixed point property of the action on the *Thurston compactification* of the Teichmüller space. Not only is this an attempt at generalization, but also it is expected to help deepen understanding of mapping classes as cluster transformations. The problem of classifying mapping classes in terms of the cluster transformations is originally raised in [PP93].

The *cluster ensemble* associated with a seed, defined in [FG09], plays a similar role of the Teichmüller space when we study cluster modular groups. It can be thought of two spaces on which the cluster modular group acts. Technically, it consists of two functors $\psi_{\mathcal{A}}, \psi_{\mathcal{X}} : \mathcal{G} \rightarrow \text{Pos}(\mathbb{R})$, called \mathcal{A} - and \mathcal{X} -spaces respectively, which are related by a natural transformation $p : \psi_{\mathcal{A}} \rightarrow \psi_{\mathcal{X}}$. Here the objects of the target category are split algebraic tori over \mathbb{R} , and the values of these functors patch together to form a pair of contractible manifolds $\mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{X}(\mathbb{R}_{>0})$, on which the cluster modular group acts analytically. These manifolds are naturally compactified to a pair of topological closed disks $\overline{\mathcal{A}} = \mathcal{A}(\mathbb{R}_{>0}) \sqcup P\mathcal{A}(\mathbb{R}^t)$ and $\overline{\mathcal{X}} = \mathcal{X}(\mathbb{R}_{>0}) \sqcup P\mathcal{X}(\mathbb{R}^t)$, called the *tropical compactifications* [FG16, Le16], on which the actions of the cluster modular group extend continuously. These are algebraic generalizations of the *Thurston compactifications* of Teichmüller spaces. In the case of the SL_2 -Teichmüller seed on a surface, $\mathcal{U}(\mathbb{R}_{>0}) = p(\mathcal{A}(\mathbb{R}_{>0}))$ is identified with the Teichmüller space, $\mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{X}(\mathbb{R}_{>0})$ are the *decorated Teichmüller space* and the *enhanced Teichmüller space* introduced by Penner [Pen87] and Fock-Goncharov [FG07], respectively. The tropical compactification $\overline{\mathcal{U}}$ is identified with the Thurston compactification of the Teichmüller space [FG16].

For each seed, a simplicial complex called the *cluster complex*, defined in [FZ03] and [FG09], admits a simplicial action of the cluster modular group. In the case of the SL_2 -Teichmüller seed on a surface F , the cluster complex is a finite covering of the *arc complex* of F . In terms of the action on the cluster complex, we define three types of elements of the cluster modular group, called *Nielsen-Thurston types*. They constitute an analogue of the classification of mapping classes.

Definition 0.1 (Nielsen-Thurston types: Definition 2.1). Let \mathbf{i} be a seed, $\mathcal{C} = \mathcal{C}_{|\mathbf{i}|}$ the corresponding cluster complex and $\Gamma = \Gamma_{|\mathbf{i}|}$ the corresponding cluster modular group. An element $\phi \in \Gamma$ is said to be

- (1) periodic if ϕ has finite order,
- (2) cluster-reducible if ϕ has a fixed point in the geometric realization $|\mathcal{C}|$ of the cluster complex, and
- (3) cluster-pseudo-Anosov (cluster-pA) if no power of ϕ is cluster-reducible.

These types give a classification of elements of the cluster modular group in the sense that the cyclic group generated by any element intersects with at least one of these types. We have the following analogue of the classical Nielsen-Thurston theory for general cluster modular groups, which is the main theorem of this paper.

Theorem 0.2 (Theorem 2.2). Let \mathbf{i} be a seed of Teichmüller type (see Definition 2.17) and $\phi \in \Gamma_{|\mathbf{i}|}$ an element. Then we have the followings.

- (1) The element $\phi \in \Gamma$ is periodic if and only if it has fixed points in $\mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{X}(\mathbb{R}_{>0})$.
- (2) The element $\phi \in \Gamma$ is cluster-reducible if and only if there exists a point $L \in \mathcal{X}(\mathbb{R}^t)_+$ such that $\phi[L] = [L]$.
- (3) If the element $\phi \in \Gamma$ is cluster-pA, there exists a point $L \in \mathcal{X}(\mathbb{R}^t) \setminus \mathcal{X}(\mathbb{R}^t)_+$ such that $\phi[L] = [L]$.

We will show that the seeds of Teichmüller type include seeds of finite type, the SL_2 -Teichmüller seeds on surfaces, and the rank 2 seeds of finite mutation type.

In the theorem above, we neither characterize cluster-pA elements in terms of fixed point property, nor describe the asymptotic behavior of the orbits as we do in the original Nielsen-Thurston classification (see Definition 3.10). However we can show the following asymptotic behavior of orbits similar to that of pA classes in the mapping class groups, for certain classes of cluster-pA elements.

Theorem 0.3 (cluster reductions and cluster Dehn twists: Theorem 2.33).

- (1) *Let \mathbf{i} be a seed, $\phi \in \Gamma_{|\mathbf{i}|}$ be a cluster-reducible element. Then some power ϕ^l induces a new element in the cluster modular group associated with a seed which has smaller mutable rank n . We call this process the cluster reduction.*
- (2) *After a finite number of cluster reductions, the element ϕ^l induces a cluster-pA element.*
- (3) *Let \mathbf{i} be a skew-symmetric connected seed which has mutable rank $n \geq 3$, $\phi \in \Gamma_{|\mathbf{i}|}$ an element of infinite order. If some power of the element ϕ is cluster-reducible to rank 2, then there exists a point $[G] \in \mathcal{PA}(\mathbb{R}^t)$ such that we have*

$$\lim_{n \rightarrow \infty} \phi^{\pm n}(g) = [G] \text{ in } \overline{\mathcal{A}}$$

for all $g \in \mathcal{A}(\mathbb{R}_{>0})$.

We call a mapping class which satisfies the assumption of Theorem 0.3(3) *cluster Dehn twist*. Dehn twists in the mapping class groups are cluster Dehn twists. The above theorem says that cluster Dehn twists have the same asymptotic behavior of orbits on $\overline{\mathcal{A}}$ as Dehn twists. We expect that cluster Dehn twists together with seed isomorphisms generate cluster modular groups, as Dehn twists do in the case of mapping class groups. The generation of cluster modular groups by cluster Dehn twists and seed isomorphisms will be discussed elsewhere.

This paper is organized as follows. In Section 1, we recall some basic definitions from [FG09]. Here we adopt slightly different treatment of the frozen vertices and definition of the cluster complex from those of [FZ03, FG09]. In Section 2, we define the Nielsen-Thurston types for elements of cluster modular groups and study the fixed point property of the actions on the tropical compactifications. Our basic examples are the SL_2 -Teichmüller seeds, studied in Section 3. Most of the contents of this section are seems to be well-known to specialists, but they are scattered in literature. Therefore we tried to gather results and give a precise description of the SL_2 -Teichmüller seeds. Other examples are studied in Appendix A.

Acknowledgement. I would like to express my gratitude to my advisor, Nariya Kawazumi, for helpful guidance and careful instruction. Also I would like to thank Toshiyuki Akita, Vladimir Fock, Rinat Kashaev, and Ken'ichi Ohshika for valuable advice and discussion. This work is partially supported by the program for Leading Graduate School, MEXT, Japan.

1. DEFINITION OF THE CLUSTER MODULAR GROUPS

1.1. The cluster modular groups and the cluster ensembles. We collect here the basic definitions on cluster ensembles and cluster modular groups. This section is based on Fock-Goncharov's seminal paper [FG09], while the treatment of frozen variables here is slightly different from them. In particular, the dimensions of the \mathcal{A} - and \mathcal{X} -spaces equal to the rank and the mutable rank of the seed, respectively. See Definition 1.9.

Definition 1.1 (seeds). A *seed* consists of the following data $\mathbf{i} = (I, I_0, \epsilon, d)$;

- (1) I is a finite set and I_0 is a subset of I called the *frozen subset*. An element of $I - I_0$ is called a *mutable vertex*.
- (2) $\epsilon = (\epsilon_{ij})$ is a \mathbb{Q} -valued function on $I \times I$ such that $\epsilon_{ij} \in \mathbb{Z}$ for $(i, j) \notin I_0 \times I_0$, which is called the *exchange matrix*.
- (3) $d = (d_i) \in \mathbb{Z}_{>0}^I$ such that $\gcd(d_i) = 1$ and the matrix $\hat{\epsilon}_{ij} := \epsilon_{ij}d_j^{-1}$ is skew-symmetric.

The seed \mathbf{i} is said to be *skew-symmetric* if $d_i = 1$ for all $i \in I$. In this case the exchange matrix ϵ is a skew-symmetric matrix. We simply write $\mathbf{i} = (I, I_0, \epsilon)$ if \mathbf{i} is skew-symmetric. We call the numbers $N := |I|$, $n := |I - I_0|$ the *rank* and the *mutable rank* of the seed \mathbf{i} , respectively.

Skew-symmetric seeds are in one-to-one correspondence with quivers without loops and 2-cycles. Here a loop is an arrow whose endpoints are the same vertex, and a 2-cycle is a pair of arrows sharing both endpoints and having different orientations. Given a skew-symmetric seed $\mathbf{i} = (I, I_0, \epsilon)$, the corresponding quiver is given by setting the set of vertices I , and drawing $|\epsilon_{ij}|$ arrows from the vertex i to the vertex j (resp. j to i) if $\epsilon_{ij} > 0$ (resp. $\epsilon_{ij} < 0$).

Definition 1.2 (seed mutations). For a seed $\mathbf{i} = (I, I_0, \epsilon, d)$ and a vertex $k \in I - I_0$, we define a new seed $\mathbf{i}' = (I', I'_0, \epsilon', d')$ as follows:

- $I' := I, I'_0 := I_0, d' := d,$
- $\epsilon'_{ij} := \begin{cases} -\epsilon_{ij} & \text{if } k \in \{i, j\}, \\ \epsilon_{ij} + \frac{|\epsilon_{ik}|\epsilon_{kj} + \epsilon_{ik}|\epsilon_{kj}|}{2} & \text{otherwise.} \end{cases}$

We write $\mathbf{i}' = \mu_k(\mathbf{i})$ and refer to this transformation of seeds as the *mutation directed to the vertex k* .

Next we associate *cluster transformation* with each seed mutation. For a field k , let k^* denote the multiplicative group. Our main interest is the case $k = \mathbb{R}$. A direct product $(k^*)^n$ is called a *split algebraic torus* over k .

Definition 1.3 (seed tori). Let $\mathbf{i} = (I, I_0, \epsilon, d)$ be a seed and $\Lambda := \mathbb{Z}[I]$, $\Lambda' := \mathbb{Z}[I - I_0]$ be the lattices generated by I and $I - I_0$, respectively.

- (1) $\mathcal{X}_{\mathbf{i}}(k) := \text{Hom}_{\mathbb{Z}}(\Lambda', k^*)$ is called the *seed \mathcal{X} -torus* associated with \mathbf{i} . For $i \in I - I_0$, the character $X_i : \mathcal{X}_{\mathbf{i}} \rightarrow k^*$ defined by $\phi \mapsto \phi(e_i)$ is called the *cluster \mathcal{X} -coordinate*, where (e_i) denotes the natural basis of Λ' .
- (2) Let $f_i := d_i^{-1}e_i^* \in \Lambda^* \otimes_{\mathbb{Z}} \mathbb{Q}$ and $\Lambda^\circ := \bigoplus_{i \in I} \mathbb{Z}f_i \subset \Lambda^* \otimes_{\mathbb{Z}} \mathbb{Q}$ another lattice, where Λ^* denotes the dual lattice of Λ and (e_i^*) denotes the dual basis of (e_i) . Then $\mathcal{A}_{\mathbf{i}}(k) := \text{Hom}_{\mathbb{Z}}(\Lambda^\circ, k^*)$ is called the *seed \mathcal{A} -torus* associated with \mathbf{i} . For $i \in I$, the character $A_i : \mathcal{A}_{\mathbf{i}} \rightarrow k^*$ defined by $\psi \mapsto \psi(f_i)$ is called the *cluster \mathcal{A} -coordinate*. The coordinates A_i ($i \in I_0$) are called *frozen variables*.

Note that $\mathcal{X}_{\mathbf{i}}(k) = (k^*)^n$ and $\mathcal{A}_{\mathbf{i}}(k) = (k^*)^N$ as split algebraic tori. These two tori are related as follows. Let $p^* : \Lambda' \rightarrow \Lambda^\circ$ be the linear map defined by

$$p^*(v) = \sum_{\substack{i \in I - I_0 \\ k \in I}} v_i \epsilon_{ik} f_k$$

for $v = \sum_{i \in I - I_0} v_i e_i \in \Lambda'$. By taking $\text{Hom}_{\mathbb{Z}}(-, k^*)$, it induces a monomial map $p_{\mathbf{i}} : \mathcal{A}_{\mathbf{i}} \rightarrow \mathcal{X}_{\mathbf{i}}$, which is represented in cluster coordinates as $p_{\mathbf{i}}^* X_i = \prod_{k \in I} A_k^{\epsilon_{ik}}$.

Remark 1.4. Note that we assign cluster \mathcal{X} -coordinates only on mutable vertices, which is a different convention from that of [FG09]. It seems to be natural to adopt our convention from the point of view of the Teichmüller theory (see Section 3).

Definition 1.5 (cluster transformations). For a mutation $\mu_k : \mathbf{i} \rightarrow \mathbf{i}'$, we define transformations on seed tori called the *cluster transformations* as follows;

- (1) $\mu_k^x : \mathcal{X}_{\mathbf{i}} \rightarrow \mathcal{X}_{\mathbf{i}'}$,
 $(\mu_k^x)^* X'_i := \begin{cases} X_k^{-1} & \text{if } i = k, \\ X_i(1 + X_k^{\text{sgn} \epsilon_{ki}})^{\epsilon_{ki}} & \text{otherwise,} \end{cases}$
- (2) $\mu_k^a : \mathcal{A}_{\mathbf{i}} \rightarrow \mathcal{A}_{\mathbf{i}'}$,
 $(\mu_k^a)^* A'_i := \begin{cases} A_i^{-1}(\prod_{\epsilon_{kj} > 0} A_j^{\epsilon_{kj}} + \prod_{\epsilon_{kj} < 0} A_j^{-\epsilon_{kj}}) & \text{if } i = k, \\ A_i & \text{otherwise.} \end{cases}$

Note that the frozen \mathcal{A} -variables are not transformed by mutations, while they have an influence on the transformations of the mutable \mathcal{A} -variables.

Definition 1.6 (the cluster modular group). Let $\mathbf{i} = (I, I_0, \epsilon, d)$ be a seed. Recall that a *groupoid* is a small category whose morphisms are all invertible.

- (1) A *seed isomorphism* is a permutation σ of I such that $\sigma(i) = i$ for all $i \in I_0$ and $\epsilon_{\sigma(i)\sigma(j)} = \epsilon_{ij}$ for all $i, j \in I$. A *seed cluster transformation* is a finite composition of mutations and seed isomorphisms. A seed cluster transformation is said to be *trivial* if the induced cluster \mathcal{A} - and \mathcal{X} -transformations are both identity. Two seeds are called *equivalent* if they are connected by a seed cluster transformation. Let $[\mathbf{i}]$ denote the equivalence class containing the seed \mathbf{i} .
- (2) Let $\mathcal{G}_{[\mathbf{i}]}$ be the groupoid whose objects are seeds in $[\mathbf{i}]$, and morphisms are seed cluster transformations, modulo trivial ones. The automorphism group $\Gamma = \Gamma_{[\mathbf{i}]} := \text{Aut}_{\mathcal{G}_{[\mathbf{i}]}}(\mathbf{i})$ is called the *cluster modular group* associated with the seed \mathbf{i} . We call elements of the cluster modular group *mapping classes* in analogy with the SL_2 -Teichmüller seeds (see Section 3).

Example 1.7. We give some examples of cluster modular groups.

- (1) (Type A_2). Let $\mathbf{i} := (\{0, 1\}, \emptyset, \epsilon)$ be the skew-symmetric seed defined by $\epsilon := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, which is called *type A_2* . Let $\phi := (0 \ 1) \circ \mu_0 \in \Gamma_{A_2}$. It is the generator of the cluster modular group. The associated cluster transformations are described as follows:

$$\begin{aligned} \phi^*(A_0, A_1) &= \left(A_1, \left(\frac{1 + A_1}{A_0} \right) \right), \\ \phi^*(X_0, X_1) &= (X_1(1 + X_0), X_0^{-1}). \end{aligned}$$

Then one can check that ϕ has order 5 by a direct calculation. See [FG09]Section 2.5 for instance. In particular we have $\Gamma_{A_2} \cong \mathbb{Z}/5$.

- (2) (Type L_k for $k \geq 2$). For an integer $k \geq 2$, let $\mathbf{i}_k := (\{0, 1\}, \emptyset, \epsilon_k)$ be the skew-symmetric seed defined by $\epsilon_k := \begin{pmatrix} 0 & k \\ -k & 0 \end{pmatrix}$. Let us refer to this seed as the *type L_k* . The quiver associated with the seed \mathbf{i}_k is shown in Figure 1. Let $\phi := (0 \ 1) \circ \mu_0 \in \Gamma_{L_k}$. It is the generator of the cluster modular group. In this case, the associated cluster transformations are described as follows:

$$\begin{aligned} \phi^*(A_0, A_1) &= \left(A_1, \left(\frac{1 + A_1^k}{A_0} \right) \right), \\ \phi^*(X_0, X_1) &= (X_1(1 + X_0)^k, X_0^{-1}). \end{aligned}$$

It turns out that in this case the element ϕ has infinite order. See Example 2.8.

$$\textcircled{0} \xrightarrow{k} \textcircled{1}$$

FIGURE 1. quiver L_k

Next we define the concept of a *cluster ensemble*, which is defined to be a pair of functors related by a natural transformation. A cluster ensemble, in particular, produces a pair of real-analytic manifolds, on which the cluster modular group acts analytically.

Let us recall some basic concepts from algebraic geometry. For a split algebraic torus H , let X_1, \dots, X_n be its coordinates. A rational function f on H is said to be *positive* if it can be represented as $f = f_1/f_2$, where $f_i = \sum_{\alpha \in \mathbb{N}^n} a_{\alpha} X^{\alpha}$ and $a_{\alpha} \in \mathbb{Z}_{\geq 0}$. Note that the set of positive rational functions on a split algebraic torus form a semifield under the usual operations. A rational map between two split algebraic tori $f : H_1 \rightarrow H_2$ is said to be *positive* if the induced map f^* preserves the semifields of positive rational functions.

Definition 1.8 (positive spaces).

- (1) Let $\text{Pos}(k)$ be the category whose objects are split algebraic tori over k and morphisms are positive rational maps. A functor $\psi : \mathcal{G} \rightarrow \text{Pos}(k)$ from a groupoid \mathcal{G} is called a *positive space*.
- (2) A *morphism* $\psi_1 \rightarrow \psi_2$ between two positive spaces $\psi_i : \mathcal{G}_i \rightarrow \text{Pos}(k)$ ($i=1,2$) consists of the data (ι, p) , where $\iota : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ is a functor and $p : \psi_1 \Rightarrow \psi_2 \circ \iota$ is a natural transformation. A morphism of positive spaces $(\iota, p) : \psi_1 \rightarrow \psi_2$ is said to be *monomial* if the map between split algebraic tori $p_{\alpha} : \psi_1(\alpha) \rightarrow \psi_2(\iota(\alpha))$ preserves the set of monomials for each object $\alpha \in \mathcal{G}_1$.

Definition 1.9 (cluster ensembles).

- (1) From Definition 1.5 we get a pair of positive spaces $\psi_{\mathcal{X}}, \psi_{\mathcal{A}} : \mathcal{G}_{|\mathbf{i}|} \rightarrow \text{Pos}(k)$, and we have a monomial morphism $p = p_{|\mathbf{i}|} : \psi_{\mathcal{A}} \rightarrow \psi_{\mathcal{X}}$ (with $\iota = \text{id}$). We call these data the *cluster ensemble* associated with the seed \mathbf{i} , and simply write as $p : \mathcal{A} \rightarrow \mathcal{X}$. The groupoid $\mathcal{G} = \mathcal{G}_{|\mathbf{i}|}$ is called the *coordinate groupoid* of the cluster ensemble.
- (2) Let $\mathcal{U} = p(\mathcal{A})$ be the positive space obtained by assigning the restriction $\psi_{\mathcal{X}}(\mu) : p_{\mathbf{i}}(\mathcal{A}_{\mathbf{i}}) \rightarrow p_{\mathbf{i}'}(\mathcal{A}_{\mathbf{i}'})$ for each mutation $\mu : \mathbf{i} \rightarrow \mathbf{i}'$.

Definition 1.10 (the positive real part). For a cluster ensemble $p : \mathcal{A} \rightarrow \mathcal{X}$ and $\mathcal{Z} = \mathcal{A}, \mathcal{U}$ or \mathcal{X} , define the *positive real part* to be the real-analytic manifold obtained by glueing seed tori by corresponding cluster transformations, as follows:

$$\mathcal{Z}(\mathbb{R}_{>0}) := \bigsqcup_{\mathbf{i} \in \mathcal{G}} \mathcal{Z}_{\mathbf{i}}(\mathbb{R}_{>0}) \Big/ (\mu_k^z),$$

where $\mathcal{Z}_{\mathbf{i}}(\mathbb{R}_{>0})$ denotes the subset of $\mathcal{Z}_{\mathbf{i}}(\mathbb{R})$ defined by the condition that all cluster coordinates are positive. Note that it is well-defined since positive rational maps preserves positive real parts. Similarly we define $\mathcal{Z}(\mathbb{Q}_{>0})$ and $\mathcal{Z}(\mathbb{Z}_{>0})$.

Note that we have a natural diffeomorphism $\mathcal{Z}_{\mathbf{i}}(\mathbb{R}_{>0}) \rightarrow \mathcal{Z}(\mathbb{R}_{>0})$ for each $\mathbf{i} \in \mathcal{G}$. The inverse map $\psi_{\mathbf{i}}^z : \mathcal{Z}(\mathbb{R}_{>0}) \rightarrow \mathcal{Z}_{\mathbf{i}}(\mathbb{R}_{>0})$ gives a chart of the manifold. The cluster modular group acts on positive real parts $\mathcal{Z}(\mathbb{R}_{>0})$ as follows:

$$(1.1) \quad \begin{array}{ccc} \mathcal{Z}(\mathbb{R}_{>0}) & \xrightarrow{\psi_{\mathbf{i}}^z} & \mathcal{Z}_{\mathbf{i}}(\mathbb{R}_{>0}) \\ \downarrow \phi & & \downarrow \mu_{i_1}^z \cdots \mu_{i_k}^z \sigma^* \\ \mathcal{Z}(\mathbb{R}_{>0}) & \xrightarrow{\psi_{\mathbf{i}'}^z} & \mathcal{Z}_{\mathbf{i}'}(\mathbb{R}_{>0}) \end{array}$$




FIGURE 2. assignment of variables

Here $\phi = \sigma \circ \mu_{i_k} \dots \mu_{i_1} \in \Gamma$ is a mapping class, σ^* is the permutation of coordinates induced by the seed isomorphism σ . The fixed point property of this action is the main subject of the present paper.

1.2. Cluster complexes. We define a simplicial complex called the *cluster complex*, on which the cluster modular group acts simplicially. In terms of the action on the cluster complex, we will define the Nielsen-Thurston types of mapping classes in Section 2. We propose here an intermediate definition between that of [FZ03] and [FG09].

Let $\mathbf{i} = (I, I_0, \epsilon, d)$ be a seed. A decorated simplex is an $(n-1)$ -dimensional simplex S with a fixed bijection, called a *decoration*, between the set of facets of S and $I - I_0$. Let \mathbb{S} be the simplicial complex obtained by glueing (infinite number of) decorated $(n-1)$ -dimensional simplices along mutable facets using the decoration. Note that the dual graph \mathbb{S}^\vee is a tree, and there is a natural covering from the set of vertices $V(\mathbb{S}^\vee)$ to the set of seeds. An edge of \mathbb{S}^\vee is projected to a mutation under this covering. Assign mutable \mathcal{A} -variables to vertices of \mathbb{S} in such a manner that:

- (1) the reflection with respect to a mutable facet takes the \mathcal{A} -variables to the \mathcal{A} -variables which are obtained by the corresponding mutation.
- (2) the labels of variables coincide with the decoration assigned to the facet in the opposite side.
- (3) the initial \mathcal{A} -coordinates are assigned to the initial simplex.

Note that the assignment is well-defined since the dual graph \mathbb{S}^\vee is a tree. Similarly we assign \mathcal{X} -variables to co-oriented facets of \mathbb{S} (see Figure 2). Let Δ be the subgroup of $\text{Aut}(\mathbb{S})$ which consists of elements that preserve all cluster variables.

Definition 1.11 (the cluster complex). The simplicial complex $\mathcal{C} = \mathcal{C}_{[\mathbf{i}]} := \mathbb{S}/\Delta$ is called the *cluster complex*. A set of vertices $\{\alpha_1, \dots, \alpha_n\} \subset V(\mathcal{C})$ is called a *cluster* if it spans a maximal simplex.

Let \mathcal{C}^\vee denote the dual graph of the cluster complex. Note that the clusters, equivalently, the vertices of \mathcal{C}^\vee , are in one-to-one correspondence with seeds together with tuples of mutable variables $((A_i), (X_i))$. For a vertex $v \in V(\mathcal{C}^\vee)$, let $[v]$ denote the underlying seed. Then we get coordinate systems of the positive real

parts for each vertex $v \in V(\mathcal{C}^\vee)$, as follows:

$$\begin{aligned}\psi_v^x : \mathcal{X}(\mathbb{R}_{>0}) &\xrightarrow{\psi_{[v]}^x} \mathcal{X}_i(\mathbb{R}_{>0}) \xrightarrow{(X_i)} \mathbb{R}_{>0}^n \\ \psi_v^a : \mathcal{A}(\mathbb{R}_{>0}) &\xrightarrow{\psi_{[v]}^a} \mathcal{A}_i(\mathbb{R}_{>0}) \xrightarrow{(A_i)} \mathbb{R}_{>0}^N\end{aligned}$$

The edges of \mathcal{C}^\vee correspond to seed mutations, and the associated coordinate transformations are described by cluster transformations.

Remark 1.12. In [FZ03], the cluster complex is defined to be a simplicial complex whose ground set is the set of mutable \mathcal{A} -coordinates, while the definition in [FG09] uses all (mutable/frozen) coordinates. In our definition, the frozen \mathcal{A} -variables have no corresponding vertices, while they are retained in the mutable \mathcal{A} -coordinates via cluster transformations.

Proposition 1.13 ([FG09]Lemma 2.15). *Let D be the subgroup of $\text{Aut}(\mathbb{S})$ which consists of elements which preserve the exchange matrix. Namely, an automorphism γ belongs to D if it satisfies $\epsilon_{\gamma(i), \gamma(j)}^{[\gamma(v)]} = \epsilon_{ij}^{[v]}$ for all $v \in V(\mathcal{C}^\vee)$ and $i, j \in [v]$. Then*

- (1) Δ is a normal subgroup of D , and
- (2) the quotient group D/Δ is naturally isomorphic to the cluster modular group Γ .

In particular, the cluster modular group acts on the cluster complex simplicially.

Example 1.14. The cluster complexes associated with seeds defined in Example 1.7 are as follows:

- (1) (Type A_2). Let \mathbf{i} be the seed of type A_2 . The cluster complex is a pentagon. The generator $\phi = (0 \ 1) \circ \mu_0 \in \Gamma_{A_2}$ acts on the pentagon by the cyclic shift.
- (2) (Type L_k for $k \geq 2$). Let \mathbf{i} be the seed of type L_k . The cluster complex is 1-dimensional, and the generator $\phi = (0 \ 1) \circ \mu_0 \in \Gamma_{L_k}$ acts by the shift of length 1. The fact that ϕ has infinite order implies that the cluster complex is the line of infinite length. See Example 2.8.

1.3. Tropical compactifications of positive spaces. Next we define tropical compactifications of positive spaces, which are described in [FG16] [Le16].

Definition 1.15 (the tropical limit). For a positive rational map $f(X_1, \dots, X_N)$ over \mathbb{R} , we define the *tropical limit* $\text{Trop}(f)$ of f by

$$\text{Trop}(f)(x_1, \dots, x_N) := \lim_{\epsilon \rightarrow 0} \epsilon \log f(e^{x_1/\epsilon}, \dots, e^{x_N/\epsilon}),$$

which defines a piecewise-linear function on \mathbb{R}^N .

Definition 1.16 (the tropical space). Let $\psi_{\mathcal{Z}} : \mathcal{G} \rightarrow \text{Pos}(\mathbb{R})$ be a positive space. Then let $\text{Trop}(\psi_{\mathcal{Z}}) : \mathcal{G} \rightarrow \text{PL}$ be the functor given by the tropical limits of positive rational maps given by $\psi_{\mathcal{Z}}$, where PL denotes the category whose objects are euclidean spaces and morphisms are piecewise-linear (PL) maps. Let $\mathcal{Z}(\mathbb{R}^t)$ be the PL manifold obtained by glueing coordinate euclidean spaces by PL maps given by $\text{Trop}(\psi_{\mathcal{Z}})$, which is called the *tropical space*.

Note that since PL maps given by tropical limits are homogeneous, $\mathbb{R}_{>0}$ naturally acts on $\mathcal{Z}(\mathbb{R}^t)$. The quotient $P\mathcal{Z}(\mathbb{R}^t) := (\mathcal{Z}(\mathbb{R}^t) \setminus \{0\})/\mathbb{R}_{>0}$ is PL homeomorphic to a sphere. Let us denote the image of $G \in \mathcal{Z}(\mathbb{R}^t) \setminus \{0\}$ under the natural projection by $[G] \in P\mathcal{Z}(\mathbb{R}^t)$. The cluster modular group acts on $\mathcal{Z}(\mathbb{R}^t)$ and $P\mathcal{Z}(\mathbb{R}^t)$ by PL homeomorphisms, similarly as eq. (1.1).

Definition 1.17 (a divergent sequence). For a positive space $\psi_{\mathcal{Z}} : \mathcal{G} \rightarrow \text{Pos}(\mathbb{R})$, we say that a sequence (g_n) in $\mathcal{Z}(\mathbb{R}_{>0})$ is *divergent* if for each compact set $K \subset \mathcal{Z}(\mathbb{R}_{>0})$ there is a number N such that $g_n \notin K$ for all $n \geq N$.

Definition 1.18 (the tropical compactification). Let $\psi_{\mathcal{X}} : \mathcal{G} \rightarrow \text{Pos}(\mathbb{R})$ be the \mathcal{X} -space associated to a seed. For a vertex $v \in V(\mathcal{C}^\vee)$, let $\mathbf{i} = [v] = (I, I_0, \epsilon, d)$ be the underlying seed, and ψ_v^x and $\text{Trop}(\psi_v^x)$ the associated positive and tropical coordinates, respectively. Then we define a homeomorphism $\mathcal{F}_v : \mathcal{X}(\mathbb{R}_{>0}) \rightarrow \mathcal{X}(\mathbb{R}^t)$ by the following commutative diagram:

$$\begin{array}{ccc} \mathcal{X}(\mathbb{R}_{>0}) & \xrightarrow{\psi_v^x} & \mathbb{R}_{>0}^n \\ \mathcal{F}_v \downarrow & & \downarrow \log \\ \mathcal{X}(\mathbb{R}^t) & \xrightarrow{\text{Trop}(\psi_v^x)} & \mathbb{R}^n \end{array} .$$

Fixing a vertex $v \in V(\mathcal{C}^\vee)$, we define the *tropical compactification* by $\overline{\mathcal{X}} := \mathcal{X}(\mathbb{R}_{>0}) \sqcup P\mathcal{X}(\mathbb{R}^t)$, and endow it with the topology of the spherical compactification. Namely, a divergent sequence (g_n) in $\mathcal{X}(\mathbb{R}_{>0})$ converges to $[G] \in P\mathcal{X}(\mathbb{R}^t)$ in $\overline{\mathcal{X}}$ if and only if $[\mathcal{F}_v(g_n)]$ converges to $[G]$ in $P\mathcal{X}(\mathbb{R}^t)$. Similarly we can consider the tropical compactifications of \mathcal{A} and \mathcal{U} -spaces, respectively.

Theorem 1.19 (Le, [Le16] Section 7). *Let $p : \mathcal{A} \rightarrow \mathcal{X}$ be a cluster ensemble, and $\mathcal{Z} = \mathcal{A}, \mathcal{U}$ or \mathcal{X} . If we have $[\mathcal{F}_v(g_n)] \rightarrow [G]$ in $P\mathcal{Z}(\mathbb{R}^t)$ for some $v \in V(\mathcal{C}^\vee)$, then we have $[\mathcal{F}_{v'}(g_n)] \rightarrow [G]$ in $P\mathcal{Z}(\mathbb{R}^t)$ for all $v' \in V(\mathcal{C}^\vee)$. In particular the definition of the tropical compactification is independent of the choice of the vertex $v \in V(\mathcal{C}^\vee)$.*

Corollary 1.20. *Let $p : \mathcal{A} \rightarrow \mathcal{X}$ be a cluster ensemble, and $\mathcal{Z} = \mathcal{A}, \mathcal{U}$ or \mathcal{X} . Then the action of the cluster modular group on the positive real part $\mathcal{Z}(\mathbb{R}_{>0})$ continuously extends to the tropical compactification $\overline{\mathcal{Z}}$.*

Proof. We need to show that $\phi_*(g_n) \rightarrow \phi_*([G])$ in $\overline{\mathcal{Z}}$ for each mapping class $\phi \in \Gamma$ and a divergent sequence (g_n) such that $g_n \rightarrow [G]$ in $\overline{\mathcal{Z}}$. Here the action in the left-hand side is given by a composition of a finite number of cluster transformations and a permutation, while the action in the right-hand side is given by its tropical limit. Then the assertion follows from Theorem 1.19. \square

Note that each tropical compactification is homeomorphic to a closed disk of an appropriate dimension.

2. NIELSEN-THURSTON TYPES ON CLUSTER MODULAR GROUPS

In this section we define three types of elements of cluster modular groups in analogy with the classical Nielsen-Thurston types (see Section 3.3). Recall that the cluster modular group acts on the cluster complex simplicially.

Definition 2.1 (Nielsen-Thurston type). Let \mathbf{i} be a seed, $\mathcal{C} = \mathcal{C}_{|\mathbf{i}|}$ be the corresponding cluster complex and $\Gamma = \Gamma_{|\mathbf{i}|}$ the corresponding cluster modular group. An element $\phi \in \Gamma$ is called

- (1) periodic if ϕ has finite order,
- (2) cluster-reducible if ϕ has a fixed point in the geometric realization $|\mathcal{C}|$ of the cluster complex, and
- (3) cluster-pseudo-Anosov (cluster-pA) if no power of ϕ is cluster-reducible.

Recall that the cluster modular group acts on the tropical compactifications $\overline{\mathcal{A}} = \mathcal{A}(\mathbb{R}_{>0}) \sqcup P\mathcal{A}(\mathbb{R}^t)$ and $\overline{\mathcal{X}} = \mathcal{X}(\mathbb{R}_{>0}) \sqcup P\mathcal{X}(\mathbb{R}^t)$, which are closed disks of dimension N and n , respectively. Hence Brouwer's fixed point theorem says that each mapping class has at least one fixed point on each of the tropical compactifications. The following is the main theorem of the present paper, which is an analogue of the classical Nielsen-Thurston classification theory.

Theorem 2.2. *Let \mathbf{i} be a seed and $\phi \in \Gamma_{|\mathbf{i}|}$ a mapping class. Then we have the followings.*

- (1) *If the mapping class $\phi \in \Gamma$ is periodic, then it has fixed points in $\mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{X}(\mathbb{R}_{>0})$.*
- (2) *If the mapping class $\phi \in \Gamma$ is cluster-reducible, then there exists a point $L \in \mathcal{X}(\mathbb{R}^t)_+$ such that $\phi[L] = [L]$.*

If the seed \mathbf{i} is of Teichmüller type (see Definition 2.17), the followings also hold:

- (1)' *if ϕ has a fixed point in $\mathcal{A}(\mathbb{R}_{>0})$ or $\mathcal{X}(\mathbb{R}_{>0})$, then ϕ is periodic.*
- (2)' *if there exists a point $L \in \mathcal{X}(\mathbb{R}^t)_+$ such that $\phi[L] = [L]$, then ϕ is cluster-reducible.*

We prove the theorem in the following several subsections. A asymptotic behavior of orbits of certain type of cluster-pA classes on the tropical compactification of the \mathcal{A} -space will be discussed in Section 2.3.

2.1. Periodic classes. Let us start by studying the fixed point property of periodic classes. Let $\mathcal{Z} = \mathcal{A}$ or \mathcal{X} .

Proposition 2.3. *Let \mathbf{i} be a seed, and $\Gamma = \Gamma_{|\mathbf{i}|}$ the associated cluster modular group. For any $\phi \in \Gamma$, consider the following conditions:*

- (i) *ϕ fixes a cell $C \in \mathcal{C}$ of finite type,*
- (ii) *ϕ is periodic, and*
- (iii) *ϕ has fixed points in $\mathcal{Z}(\mathbb{R}_{>0})$.*

Then we have (i) \Rightarrow (ii) \Rightarrow (iii). Here a cell C in the cluster complex is of finite type if the set of supercells $\{C' \in \mathcal{C} | C' > C\}$ is a finite set.

Remark 2.4. The converse assertion (iii) \Rightarrow (ii) holds under the condition (T1) on the seed. See Proposition 2.6.

Proof. (i) \Rightarrow (ii). Suppose we have $\phi(C) = C$ for some cell $C \in \mathcal{C}$ of finite type. Then from the definition, the set of supercells $\{C' \in \mathcal{C} | C' > C\}$ is a finite set, and ϕ preserves this set. Since this set contains a maximal dimensional cell and the cluster complex \mathcal{C} is connected, the action of ϕ on \mathcal{C} is determined by the action on this finite set. Hence ϕ has finite order.

(ii) \Rightarrow (iii). The proof is purely topological. Assume that ϕ has finite order. By Brouwer's fixed point theorem, ϕ has a fixed point on the disk $\overline{\mathcal{Z}} \approx D^N$. We need to show that there exists a fixed point in the interior $\mathcal{Z}(\mathbb{R}_{>0})$. Suppose ϕ has no fixed points in the interior. Then ϕ induces a homeomorphism $\tilde{\phi}$ on the sphere $S^N = D^N / \partial D^N$ obtained by collapsing the boundary to a point, and $\tilde{\phi}$ has no fixed points other than the point corresponding to the image of ∂D^N . Now we use the following theorem.

Theorem 2.5 (Brown [Bro82] Theorem 5.1). *Let X be a paracompact space of finite cohomological dimension, s a homeomorphism of X , which has finite order. If $H_*(\text{Fix}(s^k); \mathbb{Z})$ is finitely generated for each k , then the Lefschetz number of s equals the Euler characteristic of the fixed point set;*

$$\text{Lef}(s) := \sum_i \text{Tr}(s : H_i(X) \rightarrow H_i(X)) = \chi(\text{Fix}(s)).$$

Applying Brown's theorem for $X = S^N$ and $s = \tilde{\phi}$ we get a contradiction, since the Lefschetz number of $\tilde{\phi}$ is an even number in this case, while the Euler characteristic of a point is 1. Hence ϕ has a fixed point in the interior $\mathcal{Z}(\mathbb{R}_{>0})$. \square

To get the converse implication (iii) \Rightarrow (ii), we need a condition on the seed, which can be thought of an algebraic formulation of the proper discontinuity of the action of the cluster modular group on positive spaces.

Proposition 2.6 (Growth property (T1)). *Suppose that a seed \mathbf{i} satisfies the following condition.*

- (T1) *For any vertex $v_0 \in V(\mathcal{C}^\vee)$, $g \in \mathcal{Z}(\mathbb{R}_{>0})$ and a number $M > 0$, there exists a number $B > 0$ such that $\max_{\alpha \in v} |\log Z_\alpha(g)| \geq M$ for all vertex $v \in V(\mathcal{C}^\vee)$ such that $[v] = [v_0]$ and $d_{\mathcal{C}^\vee}(v, v_0) \geq B$.*

Then the conditions (ii) and (iii) in Proposition 2.3 are equivalent. Here $d_{\mathcal{C}^\vee}$ denotes the graph metric on the 1-skelton of \mathcal{C}^\vee .

Roughly speaking, the condition (T1) says that the values of the cluster coordinates evaluated at a point g diverge as we perform a sequence of mutations which increase the distance $d_{\mathcal{C}^\vee}$.

Proof. Let $\phi \in \Gamma_{|\mathbf{i}|}$ be an element of infinite order. We need to show that ϕ has no fixed points in $\mathcal{Z}(\mathbb{R}_{>0})$. It suffices to show that each orbit is divergent. Let $g \in \mathcal{Z}(\mathbb{R}_{>0})$ and $K \subset \mathcal{Z}(\mathbb{R}_{>0})$ a compact set. We claim that there exists a number N such that $\phi^n(g) \notin K$ for all $n \geq N$. Take a number $M > 0$ so that $M \geq \max_{i=1, \dots, N} \max_{g \in K} |\log Z_i(g)|$.

Note that since the 1-skelton of \mathcal{C}^\vee has valency n at any vertex, the graph metric $d_{\mathcal{C}^\vee}$ is proper. Namely, the number of vertices v such that $d_{\mathcal{C}^\vee}(v, v_0) \leq B$ is finite for any $B > 0$. Hence for the number $B > 0$ given by the assumption (T1), there exists a number N such that $d_{\mathcal{C}^\vee}(\phi^{-n}(v_0), v_0) \geq B$ for all $n \geq N$, since ϕ has infinite order. Also note that $[\phi^{-n}(v_0)] = [v_0]$ by Proposition 1.13. Then we have

$$\max_{i=1, \dots, N} |\log Z_i(\phi^n(g))| = \max_{\alpha \in \phi^{-n}(v_0)} |\log Z_\alpha(g)| \geq M$$

for all $n \geq N$, where (Z_1, \dots, Z_n) is the coordinate system associated with the vertex v_0 . Here we used the equivariance of the coordinates $Z_{\phi^{-1}(\alpha)}(g) = Z_\alpha(\phi(g))$. Hence we have $\phi^n(g) \notin K$ for all $n \geq N$. \square

Proposition 2.7. *Assume that the cluster modular group $\Gamma_{|\mathbf{i}|}$ acts on $\mathcal{Z}_{|\mathbf{i}|}(\mathbb{R}_{>0})$ proper discontinuously. Then the condition (T1) holds.*

Proof. Suppose that the condition (T1) does not hold. Then there exists a vertex $v_0 \in V(\mathcal{C}^\vee)$, a point $g \in \mathcal{Z}(\mathbb{R}_{>0})$, a number $M > 0$, and a sequence $(v_n) \subset V(\mathcal{C}^\vee)$ such that $d_{\mathcal{C}^\vee}(v_n, v_0) \geq n$ and $\max_{\alpha \in v_n} |\log Z_\alpha(g)| \leq M$. Take a mapping class $\psi_n \in \Gamma$ so that $\psi_n(v_n) = v_0$. It is possible since $[v_n] = [v_0]$. Then we have

$$\max_{i=1, \dots, N} |\log Z_i(\psi_n(g))| = \max_{\alpha \in \psi_n^{-1}(v_n)} |\log Z_\alpha(g)| \leq M,$$

which implies that there exists a compact set $K \subset \mathcal{Z}(\mathbb{R}_{>0})$ such that $\psi_n(g) \in K$ for all n . Note that the mapping classes (ψ_n) are distinct, since the vertices (v_n) are distinct. In particular we have $\psi_n^{-1}(K) \cap K \neq \emptyset$ for all n , consequently the action is not properly discontinuous. \square

We will verify the condition (T1) for SL_2 -Teichmüller seeds using Proposition 2.7 in Section 3.2, and for the simplest case L_k ($k \geq 2$) of infinite type in Appendix A.

Example 2.8.

- (1) (Type A_2). Let \mathbf{i} be the seed of type A_2 and $\phi = (0 \ 1) \circ \mu_0 \in \Gamma_{A_2}$ the generator. See Example 1.7. Recall that the two actions on the positive spaces $\mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{X}(\mathbb{R}_{>0})$ are described as follows:

$$\begin{aligned} \phi^*(A_0, A_1) &= \left(A_1, \left(\frac{1 + A_1}{A_0} \right) \right), \\ \phi^*(X_0, X_1) &= (X_1(1 + X_0), X_0^{-1}). \end{aligned}$$

The fixed points are given by $(A_0, A_1) = ((1 + \sqrt{5})/2, (1 + \sqrt{5})/2)$ and $(X_0, X_1) = ((1 + \sqrt{5})/2, (-1 + \sqrt{5})/2)$, respectively.

- (2) (Type L_k for $k \geq 2$). Let \mathbf{i} be the seed of type L_k and $\phi = (0 \ 1) \circ \mu_0 \in \Gamma_{L_k}$ the generator. See Example 1.7. Recall that the two actions on the positive spaces $\mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{X}(\mathbb{R}_{>0})$ are described as follows:

$$\begin{aligned}\phi^*(A_0, A_1) &= \left(\frac{1 + A_1^k}{A_0} \right), \\ \phi^*(X_0, X_1) &= (X_1(1 + X_0)^k, X_0^{-1}).\end{aligned}$$

These equations have no positive solutions. Indeed, the \mathcal{X} -equation implies $X_0^2 = (1 + X_0)^k$, which has no positive solution since ${}_kC_2 \geq 1$ for $k \geq 2$. Similarly for \mathcal{A} -variables. Hence we can conclude that ϕ has infinite order by Proposition 2.3. In particular we have $\Gamma_{L_k} \cong \mathbb{Z}$.

2.2. Cluster-reducible classes. In this subsection, we study the fixed point property of a cluster-reducible class. The basic idea is the following. For the SL_2 -Teichmüller seed on a marked hyperbolic surface F of type F_g^1 or $F_{g,\delta}$, the vertices of the cluster complex \mathcal{C} are *ideal arcs* on F . See Theorem 3.6. In particular each point in the geometric realization $|\mathcal{C}|$ of the cluster complex is represented by the projective class of a linear combination of some ideal arcs. On the other hand, the projectivized tropical \mathcal{X} -space $P\mathcal{X}(\mathbb{R}^t)$ contains all such projective classes. Hence the cluster complex is embedded into the projectivized tropical \mathcal{X} -space in this case. In Section 2.2.1 we show that this picture is valid for general seeds, under some condition on the seed. See Lemma 2.11.

2.2.1. Fixed points in the tropical \mathcal{X} -space.

Definition 2.9 (the non-negative part). Let \mathbf{i} be a seed. For any vertex $v \in V(\mathcal{C}^\vee)$, let $C_v := \{L \in \mathcal{X}(\mathbb{R}^t) \mid L \geq 0 \text{ in } v\}$ be a cone in the tropical space, where $L \geq 0$ in v means that $x_\alpha(L) \geq 0$ for all $\alpha \in v$. Then the union $\mathcal{X}(\mathbb{R}^t)_+ := \bigcup_{v \in V(\mathcal{C}^\vee)} C_v \subseteq \mathcal{X}(\mathbb{R}^t)$ is called the *non-negative part* of $\mathcal{X}(\mathbb{R}^t)$.

Let us define a Γ -equivariant map $\Psi : \mathcal{C} \rightarrow P\mathcal{X}(\mathbb{R}^t)_+$ as follows. The construction contains reformulations of some conjectures stated in [FG09] Section 5, for later use. For each maximal simplex S of \mathbb{S} , let $v \in V(\mathcal{C}^\vee)$ be the vertex corresponding to the maximal simplex given by the image of S in \mathcal{C} . Then we have the following map:

$$\Psi_S : S \cong P\mathbb{R}_{\geq 0}^n \xrightarrow{\xi_v^{-1}} PC_v \subseteq P\mathcal{X}(\mathbb{R}^t)_+,$$

where $\xi_v := \text{Trop}(\psi_v^x) : \mathcal{X}(\mathbb{R}^t) \rightarrow \mathbb{R}^n$ is the tropical coordinate associated with the vertex v , whose restriction gives a bijection $C_v \rightarrow \mathbb{R}_{\geq 0}^n$. Since the tropical \mathcal{X} -transformation associated to a mutation $\mu_k : v \rightarrow v'$ preserves the set $\{x_k = 0\}$ and the dual graph \mathbb{S}^\vee is a tree, these maps combine to give a map

$$\Psi := \bigcup_{v \in V(\mathcal{C}^\vee)} \Psi_v : \mathbb{S} \rightarrow P\mathcal{X}(\mathbb{R}^t)_+,$$

which is clearly surjective. Assume we have $S' = \gamma(S)$ for some $\gamma \in \Delta$. Then from the definition of Δ , γ preserves all the tropical \mathcal{X} -coordinates. Hence we have $\Psi_{v'}(\gamma x) = \Psi_v(x)$ for all $x \in S$, and the map descends to

$$\Psi : \mathcal{C} = \mathbb{S}/\Delta \rightarrow P\mathcal{X}(\mathbb{R}^t)_+.$$

Lemma 2.10. *The surjective map Ψ defined above is Γ -equivariant.*

Proof. It follows from the following commutative diagram for $\phi \in \Gamma$:

$$\begin{array}{ccccc} [S_v] & \xrightarrow{=} & P\mathbb{R}_{\geq 0}^n & \xrightarrow{\xi_v^{-1}} & C_v \\ \downarrow \phi & & \downarrow \phi^* & & \downarrow \phi^x \\ [S_{\phi^{-1}(v)}] & \xrightarrow{=} & P\mathbb{R}_{\geq 0}^n & \xrightarrow{\xi_{\phi^{-1}(v)}^{-1}} & PC_v \end{array}$$

Here $[S_v]$ is the cluster in \mathcal{C} which corresponds to the vertex $v \in V(\mathcal{C}^\vee)$, ϕ^* is the permutation on vertices induced by ϕ , and ϕ^x is the induced tropical \mathcal{X} -transformation on $\mathcal{X}(\mathbb{R}^t)$. \square

Next we introduce a sufficient condition for Ψ being injective. For a point $L \in \mathcal{X}(\mathbb{R}^t)$, a seed \mathbf{i} is called a *non-negative seed* for L if $L \in C_{\mathbf{i}}$. The set of vertices whose coordinates of L are zero defines a subseed $\mathbf{j}(L) \subset \mathbf{i}$, called the *zero subseed* of L . Since the mutation directed to a vertex $k \in \mathbf{j}(L)$ preserves the sign of coordinates, there is a unique subseed of $\mu_k(\mathbf{i})$ that corresponds to $\mathbf{j}(L)$. Two non-negative seeds \mathbf{i} and \mathbf{i}' are called *$\mathbf{j}(L)$ -equivalent* if they are connected by a finite sequence of mutations directed to the vertices in $\mathbf{j}(L)$.

Lemma 2.11. *Assume that a seed \mathbf{i} satisfies the following condition:*

(T2) *For each $L \in \mathcal{X}(\mathbb{R}^t)_+$, any two non-negative seeds for L are $\mathbf{j}(L)$ -equivalent.*

Then the map $\Psi : \mathcal{C} \rightarrow P\mathcal{X}(\mathbb{R}^t)_+$ is a Γ -equivariant isomorphism.

Compare the condition (T2) with Conjecture 5.10 in [FG09].

Proof. We need to prove the injectivity of Ψ . Note that Ψ is injective on each simplex.

Assume that $x \in [S_{\mathbf{i}}], x' \in [S_{\mathbf{i}'}]$ for $\mathbf{i} \neq \mathbf{i}'$ and $\Psi(x) = \Psi(x') =: [L] \in P\mathcal{X}(\mathbb{R}^t)_+$. Let $\mathbf{j}(L) \subset \mathbf{i}$ be the zero subseed. If $x \in \text{int}[S_{\mathbf{i}}]$, then $L > 0$ in \mathbf{i} and $\mathbf{j}(L) = \emptyset$. Hence from the first paragraph, we have $x = x'$. If $\mathbf{j}(L) \neq \emptyset$, then by assumption \mathbf{i}' is $\mathbf{j}(L)$ -equivalent to \mathbf{i} . Then $x, x' \in [S_{\mathbf{j}(L)}] \subset [S_{\mathbf{i}}] \cap [S_{\mathbf{i}'}]$. Hence from the first paragraph, we have $x = x'$. Therefore Ψ is injective. \square

Example 2.12. Seeds of finite type satisfy the equivalence property (T2), see [FG09] Theorem 5.8.

Proposition 2.13 (fixed points in \mathcal{X} -space). *Let \mathbf{i} be a seed, and $\phi \in \Gamma_{|\mathbf{i}|}$ a mapping class. Then we have the followings.*

- (1) *If ϕ is cluster-reducible, then there is a point $L \in \mathcal{X}(\mathbb{R}^t)_+ \setminus \{0\}$ such that $\phi[L] = [L]$.*
- (2) *If \mathbf{i} satisfies the condition (T2), then the converse of (1) is also true.*

Proof. The assertions follow from Lemma 2.10 and Lemma 2.11, respectively. \square

Definition 2.14 (seeds of definite type). A seed \mathbf{i} is of *definite type* if $\mathcal{X}_{|\mathbf{i}|}(\mathbb{R}^t)_+ = \mathcal{X}_{|\mathbf{i}|}(\mathbb{R}^t)$.

Proposition 2.15. *Assume that a seed \mathbf{i} satisfies the equivalence property (T2). Then \mathbf{i} is of definite type if and only if it is of finite type.*

Proof. The fact that finite type seeds are definite is due to Fock-Goncharov [FG09]. Let us prove the converse implication. Assume that \mathcal{X} is of definite type. Then by Lemma 2.11 we have a homeomorphism $\Psi : \mathcal{C} \rightarrow P\mathcal{X}(\mathbb{R}^t)$, and the latter is homeomorphic to a sphere. In particular \mathcal{C} is a compact simplicial complex, hence it can possess finitely many cells. \square

Remark 2.16. The conclusion part in Proposition 2.15 is Conjecture 5.7 in [FG09].

Definition 2.17 (seeds of Teichmüller type). A seed \mathbf{i} is of Teichmüller type if it satisfies the growth property (T1) and equivalence property (T2), defined in Proposition 2.6 and Lemma 2.11, respectively.

Example 2.18.

- (1) Seeds of finite type are of Teichmüller type. See [FG09] Theorem 5.8.
- (2) SL_2 -Teichmüller seeds are of Teichmüller type. See Section 3.2.
- (3) The seed of type L_k ($k \geq 1$) is of Teichmüller type. See Appendix A.

Corollary 2.19. *Let \mathbf{i} be a seed of Teichmüller type, and $\phi \in \Gamma$ a cluster- $p\mathcal{A}$ class. Then there exists a point $L \in \mathcal{X}(\mathbb{R}^t) \setminus \mathcal{X}(\mathbb{R}^t)_+$ such that $\phi[L] = [L]$.*

Proof. Since the tropical compactification $\overline{\mathcal{X}}$ is a closed disk, Brouwer's fixed point theorem says that there exists a point $x \in \overline{\mathcal{X}}$ such that $\phi(x) = x$. If $x \in \mathcal{X}(\mathbb{R}_{>0})$, then by assumption ϕ has finite order, which is a contradiction. If $x \in P\mathcal{X}(\mathbb{R}^t)_+$, then by Proposition 2.13(2), ϕ is cluster-reducible, which is a contradiction. Hence $x \in P(\mathcal{X}(\mathbb{R}^t) \setminus \mathcal{X}(\mathbb{R}^t)_+)$. \square

2.2.2. *cluster reduction and fixed points in the tropical \mathcal{A} -space.* Here we define an operation, called the *cluster reduction*, which produces a new seed from a given seed and a certain set of vertices of the cluster complex. In the last of this subsection we study the fixed point property of a cluster-reducible class on the tropical \mathcal{A} -space.

Let $\{\alpha_1, \dots, \alpha_k\} \subset V(\mathcal{C})$ be a subset of vertices, which is contained in a cluster.

Lemma 2.20 (the cluster reduction of a seed). *Take a cluster containing $\{\alpha_1, \dots, \alpha_k\}$. Let $\mathbf{i} = (I, I_0, \epsilon, d)$ be the underlying seed and $i_j := [\alpha_j] \in I$ the corresponding vertex for $j = 1, \dots, n-2$ under the projection $[\] : \{\text{clusters}\} \rightarrow \{\text{seeds}\}$ (see Definition 1.11). Then we define a new seed by $\mathbf{i}' := (I, I_0 \sqcup \{i_1, \dots, i_k\}, \epsilon, d)$, namely, by "freezing" the vertices $\{i_1, \dots, i_k\}$. Then the corresponding cluster complex $\mathcal{C}' := \mathcal{C}_{|\mathbf{i}'|}$ is naturally identified with the link of $\{\alpha_1, \dots, \alpha_k\}$ in \mathcal{C} . In particular the equivalence class $|\mathbf{i}'|$ does not depend on the choice of the cluster containing $\{\alpha_1, \dots, \alpha_k\}$.*


Proof. Let $C \subset \mathcal{C}$ be a cluster containing $\{\alpha_1, \dots, \alpha_k\}$, $\mathbf{i} = (I, I_0, \epsilon, d)$ the corresponding seed. For a mutation directed to a mutable vertex $k \in I - (I_0 \sqcup \{i_1, \dots, i_k\})$, the cluster $C' = \mu_k(C)$ also contains $\{\alpha_1, \dots, \alpha_k\}$. Conversely, any cluster C' containing $\{\alpha_1, \dots, \alpha_k\}$ is obtained by such a sequence of mutations. Hence each cluster in the cluster complex \mathcal{C}' has the form $C \setminus \{\alpha_1, \dots, \alpha_k\}$, for some cluster $C \subset \mathcal{C}$ containing $\{\alpha_1, \dots, \alpha_k\}$. \square

We say that the corresponding object, such as the cluster ensemble $p_{|\mathbf{i}'|} : \mathcal{A}_{|\mathbf{i}'|} \rightarrow \mathcal{X}_{|\mathbf{i}'|}$ or the cluster modular group $\Gamma_{|\mathbf{i}'|}$, is obtained by the *cluster reduction* with respect to the invariant set $\{\alpha_1, \dots, \alpha_k\}$ from the original one. Next we show that some power of a cluster-reducible class induces a new mapping class by the cluster reduction.

Lemma 2.21. *Let \mathbf{i} be a seed, $\phi \in \Gamma_{|\mathbf{i}|}$ a mapping class. Then ϕ is cluster-reducible if and only if it has an invariant set of vertices $\{\alpha_1, \dots, \alpha_k\} \in V(\mathcal{C})$ contained in a cluster.*

Proof. Suppose ϕ is cluster-reducible. Then ϕ has a fixed point $c \in |\mathcal{C}|$. Since the action is simplicial, ϕ fixes the cluster C containing the point c . Hence ϕ permutes the vertices of C , which give an invariant set contained in C . The converse is also true, since ϕ fixes the point given by the barycenter of the vertices $\{\alpha_1, \dots, \alpha_k\}$. \square

Definition 2.22 (proper reducible classes). A mapping class $\phi \in \Gamma_{|\mathbf{i}|}$ is called *proper reducible* if it has a fixed point in $V(\mathcal{C})$.

FIGURE 3. quiver X_7

Lemma 2.23. *Let $\phi \in \Gamma_{|\mathbf{i}|}$ be a mapping class.*

- (1) *If ϕ is proper reducible, then ϕ is cluster reducible.*
- (2) *If ϕ is cluster-reducible, then some power of ϕ is proper reducible.*

Proof. Clear from the previous lemma. \square

Lemma 2.24 (the cluster reduction of a proper reducible class). *Let $\phi \in \Gamma_{|\mathbf{i}|}$ be a proper reducible class, $\{\alpha_1, \dots, \alpha_k\}$ a fixed point set of vertices contained in a cluster. Then ϕ induces a new mapping class $\phi' \in \Gamma_{|\mathbf{i}'|}$ in the cluster modular group obtained by the cluster reduction with respect to $\{\alpha_1, \dots, \alpha_k\}$.*

Proof. The identification of \mathcal{C}' with the link of the invariant set $\{\alpha_1, \dots, \alpha_k\}$ in \mathcal{C} induces an group isomorphism

$$\Gamma_{|\mathbf{i}'|} \cong \{\psi \in \Gamma_{|\mathbf{i}|} \mid \psi(\mathcal{C}') = \mathcal{C}'\},$$

and the right-hand side contains ϕ . Let $\phi' \in \Gamma_{|\mathbf{i}'|}$ be the corresponding element. Note that ϕ fixes all frozen vertices in \mathbf{i}' , since it is proper reducible. \square

We say that the mapping class ϕ' is obtained by the *cluster reduction* with respect to the fixed point set $\{\alpha_1, \dots, \alpha_k\}$ from ϕ .

Lemma 2.25. *A proper reducible class of infinite order induces a cluster-pA class in the cluster modular group corresponding to the seed obtained by a finite number of the cluster reductions.*

Proof. Clear from the definition of the cluster-pA classes. \square

Example 2.26 (Type X_7). Let $\mathbf{i} = (\{0, 1, 2, 3, 4, 5, 6\}, \emptyset, \epsilon)$ be the skew-symmetric seed defined by the quiver described in Figure 3. We call this seed *type X_7* , following [FST12]. The mapping class $\phi_1 := (1\ 2) \circ \mu_1 \in \Gamma_{X_7}$ is proper reducible and fixes the vertex $A_i \in V(\mathcal{C})$ ($i = 0, 3, 4, 5, 6$), which is the i -th coordinate in the initial cluster. The cluster reduction with respect to the invariant set $\{A_0, A_3, A_4, A_5, A_6\}$ produces a seed $\mathbf{i}' = (\{0, 1, 2, 3, 4, 5, 6\}, \{0, 3, 4, 5, 6\}, \epsilon)$ of type L_2 , except for some non-trivial coefficients. The cluster complex $\mathcal{C}_{|\mathbf{i}'|}$ is identified with the link of $\{A_0, A_3, A_4, A_5, A_6\}$, which is the line of infinite length. The cluster reduction ϕ' is cluster-pA, and acts on this line by the shift of length 1. Compare with Example 1.14.

The mapping class $\psi_1 := (0\ 1\ 2)(3\ 4\ 5\ 6) \circ \mu_2 \mu_1 \mu_0 \in \Gamma_{X_7}$ is cluster-reducible, since it has an invariant set $\{A_3, A_4, A_5, A_6\}$ contained in the initial cluster. Note that the power ψ_1^2 is proper reducible, since it fixes the vertex A_0 .

Lemma 2.27. *Let \mathbf{i} be a seed, and \mathbf{i}' the seed obtained by a cluster reduction. Let $\psi_{\mathcal{A}} : \mathcal{G}_{|\mathbf{i}|} \rightarrow \text{Pos}(\mathbb{R})$ and $\psi'_{\mathcal{A}} : \mathcal{G}_{|\mathbf{i}'|} \rightarrow \text{Pos}(\mathbb{R})$ be the positive \mathcal{A} -spaces associated with the seeds \mathbf{i} and \mathbf{i}' , respectively. Then there is a natural morphism of the positive spaces $(\iota, q) : \psi'_{\mathcal{A}} \rightarrow \psi_{\mathcal{A}}$ which induces $\Gamma_{|\mathbf{i}'|}$ -equivariant homeomorphisms $\mathcal{A}'(\mathbb{R}_{>0}) \cong \mathcal{A}(\mathbb{R}_{>0})$ and $\mathcal{A}'(\mathbb{R}^t) \cong \mathcal{A}(\mathbb{R}^t)$.*

Proof. Note that the only difference between the two positive \mathcal{A} -spaces is the admissible directions of mutations. The functor $\iota : \mathcal{G}_{|\mathbf{i}'|} \rightarrow \mathcal{G}_{|\mathbf{i}|}$ between the coordinate groupoids is defined by $(I, I_0 \sqcup \{i_1, \dots, i_k\}, \epsilon, d) \mapsto (I, I_0, \epsilon, d)$ and sending the morphisms naturally. The identity map $\mathcal{A}_{\mathbf{i}'}(k) = \mathcal{A}_{\mathbf{i}}(k)$ for each \mathcal{A} -torus combine to give a natural transformation $q : \psi'_{\mathcal{A}} \Rightarrow \psi_{\mathcal{A}} \circ \iota$. The latter assertion is clear. \square

Remark 2.28. We have no natural embedding of the \mathcal{X} -space in general, since \mathcal{X} -coordinates assigned to the vertices in $\{i_1, \dots, i_k\}$ may be changed by cluster \mathcal{X} -transformations directed to the vertices in $I - (I_0 \sqcup \{i_1, \dots, i_k\})$.

Definition 2.29. A tropical point $G \in \mathcal{A}(\mathbb{R}^t)$ is said to be *cluster-filling* if it satisfies $a_{\alpha}(G) \neq 0$ for all $\alpha \in V(\mathcal{C})$.

Note that the definition depends only on the projective class of G .

Proposition 2.30 (fixed points in \mathcal{A} -space). *Let \mathbf{i} be a seed satisfying the condition (T1), and $\Gamma = \Gamma_{|\mathbf{i}|}$ the corresponding cluster modular group. For a proper reducible class $\phi \in \Gamma$ of infinite order, there exists a non-cluster-filling point $G \in \mathcal{A}(\mathbb{R}^t)$ such that $\phi[G] = [G]$.*

Proof. Let $\{\alpha_1, \dots, \alpha_k\}$ be a fixed point set of ϕ contained in a cluster, and $\phi' \in \Gamma_{|\mathbf{i}'|}$ the corresponding cluster reduction. Since the tropical compactification $\overline{\mathcal{A}'}$ is a closed disk, ϕ' has a fixed point $x' \in \overline{\mathcal{A}'}$ by Brouwer's fixed point theorem. By Proposition 2.6, x must be a point on the boundary $P\mathcal{A}'(\mathbb{R}^t)$. Then ϕ fixes the image x of $x' \in \overline{\mathcal{A}}$ under the homeomorphism given by Lemma 2.27. \square

2.3. Cluster-pA classes of special type: cluster Dehn twists. Using the cluster reduction we define special type of cluster-pA mapping classes, called *cluster Dehn twists*, and prove that they have an asymptotic behavior of orbits on the tropical compactification of the \mathcal{A} -space analogous to that of Dehn twists in the mapping class group.

Definition 2.31 (cluster Dehn twists). Let \mathbf{i} be a skew-symmetric seed of mutable rank n . A cluster-reducible class $\phi \in \Gamma_{|\mathbf{i}|}$ is said to be *cluster-reducible to rank m* if the following conditions hold.

- (1) There exists a number $l \in \mathbb{Z}$ such that $\psi = \phi^l$ is proper reducible.
- (2) The mapping class ψ induces a mapping class in the cluster modular group associated with the seed of mutable rank m obtained by the cluster reduction with respect to a fixed point set $\{\alpha_1, \dots, \alpha_{n-m}\}$ of ψ .

A cluster-reducible class ϕ of infinite order is called a *cluster Dehn twist* if it is cluster-reducible to rank 2. Namely, there exists a number $l \in \mathbb{Z}$ and a subset $\{\alpha_1, \dots, \alpha_{n-2}\} \subset V(\mathcal{C}_{|\mathbf{i}|})$ of vertices which is fixed by ϕ^l and contained in a cluster, where n is the mutable rank of \mathbf{i} .

A skew-symmetric seed is said to be *connected* if the corresponding quiver is connected.

Lemma 2.32. *Let \mathbf{i} be a skew-symmetric connected seed of mutable rank $n \geq 3$. Suppose that a proper reducible class $\psi \in \Gamma_{|\mathbf{i}|}$ has infinite order and there exists a subset $\{\alpha_1, \dots, \alpha_{n-2}\} \subset V(\mathcal{C}_{|\mathbf{i}|})$ of vertices which is fixed by ψ and contained in*

a cluster. Then the action of the cluster reduction $\psi' \in \Gamma_{|\mathbf{i}'|}$ with respect to the invariant set $\{\alpha_1, \dots, \alpha_{n-2}\}$ on the \mathcal{A} -space is represented as follows:

$$(2.1) \quad (\psi')^*(A_0, A_1) = \left(A_1, \frac{C + A_1^2}{A_0} \right).$$

Here (A_0, A_1) denotes the remaining cluster coordinates of the \mathcal{A} -space under the cluster reduction, C is a product of frozen variables.

Proof. Take a cluster containing $\{\alpha_1, \dots, \alpha_{n-2}\}$. Let $\mathbf{i} = (I, I_0, \epsilon)$ be the corresponding seed, and $i_j := [\alpha_j] \in I$ the corresponding vertex for $j = 1, \dots, n-2$. Then the cluster reduction produces a new seed $\mathbf{i}' := (I, I_0 \sqcup \{i_1, \dots, i_{n-2}\}, \epsilon)$, whose mutable rank is 2. Label the vertices so that $I - I'_0 = \{0, 1\}$ and $I'_0 = \{2, \dots, N-1\}$, where $I'_0 := I_0 \sqcup \{i_1, \dots, i_{n-2}\}$ and N is the rank of the seed \mathbf{i} . Note that $\psi = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \circ \mu_0 \in \Gamma_{|\mathbf{i}|}$. Suitably relabeling if necessary, we can assume that $k := \epsilon_{01} > 0$. We claim that $k = 2$. Since \mathbf{i} is connected, there exists a vertex $i \in I'_0$ such that $a := \epsilon_{i0} \neq 0$ or $b := \epsilon_{1i} \neq 0$. Since ψ preserves the quiver, we compute that $a = b$ and $b - ak = -a$. Hence we conclude that $k = 2$. Then from the definition of the cluster \mathcal{A} -transformation we have

$$\psi^*(A_0, A_1) = \left(A_1, \frac{\prod_{i \in I'_0} A_i^{\epsilon_{i0}} + A_1^2}{A_0} \right)$$

and $\psi^*(A_i) = A_i$ for all $i \in I'_0$, as desired. \square

Theorem 2.33. *Let \mathbf{i} be a skew-symmetric connected seed of mutable rank $n \geq 3$ or the seed of type L_2 . Then for each cluster Dehn twist $\phi \in \Gamma_{|\mathbf{i}|}$, there exists a cluster-filling point $[G] \in P\mathcal{A}(\mathbb{R}^t)$ such that we have*

$$\lim_{n \rightarrow \infty} \phi^{\pm n}(g) = [G] \text{ in } \overline{\mathcal{A}}$$

for all $g \in \mathcal{A}(\mathbb{R}_{>0})$.


Proof. Assume that $n \geq 3$. There exists a number l such that $\psi := \phi^l$ satisfies the assumption of Lemma 2.32. Let us consider the following recurrence relation:

$$\begin{cases} a_0^{(n)} &= -a_1^{(n-1)}, \\ a_1^{(n)} &= -a_0^{(n-1)} + \log(C + e^{2a_1^{(n-1)}}), \end{cases}$$

where $C > 0$ is a positive constant. It is the log-dynamics of eq. (2.1). Then one can directly compute that $a_0^{(n)}, a_1^{(n)}$ goes to infinity and $a_0^{(n)}/a_1^{(n)} \rightarrow 1$ as $n \rightarrow \infty$ for arbitrary initial real values. Hence we conclude that $\psi^n(g) \rightarrow [G]$ in $\overline{\mathcal{A}}$ for all g in $\mathcal{A}(\mathbb{R}_{>0})$, where $G \in \mathcal{A}(\mathbb{R}^t)$ is the point whose coordinates are $a_0 = a_1 = 1$, $a_i = 0$ for all $i \in I'_0$. The proof for the negative direction is similar. The generator of Γ_{L_2} , which is cluster-pA, also satisfies the desired property. \square

Example 2.34 (Dehn twists in the mapping class group). Let $F = F_g^s$ be a hyperbolic surface with $s \geq 2$. For an essential non-separating simple closed curve C , we denote the right hand Dehn twist along C by $t_C \in MC(F)$. Consider an annular neighborhood $\mathcal{N}(C)$ of C , and slide two of punctures so that exactly one puncture lies on each boundary component of $\mathcal{N}(C)$. Let Δ be an ideal triangulation obtained by glueing the ideal triangulation of $\mathcal{N}(C)$ shown in Figure 4 and an arbitrary ideal triangulation of $F \setminus \mathcal{N}(C)$. This kind of a triangulation is given in [Kas01]. Then the Dehn twist is represented as $t_C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \circ \mu_0$, hence it is a cluster Dehn twist with $l = 1$. Its action on the \mathcal{A} -space is represented as

$$\phi_1^*(A_0, A_1, A_2, A_3) = \left(A_1, \frac{A_2 A_3 + A_1^2}{A_0}, A_2, A_3 \right).$$

FIGURE 4. ideal triangulation of $\mathcal{N}(C)$

Example 2.35 (Type X_7). Let us consider the seed of *type* X_7 . The mapping class $\phi_1 := (1\ 2) \circ \mu_1 \in \Gamma_{X_7}$ is a cluster Dehn twist, whose action on the \mathcal{A} -space is represented as

$$\phi_1^*(A_0, A_1, A_2) = \left(A_0, A_2, \frac{A_0 + A_2^2}{A_1} \right).$$

For a general cluster-pA class, we only know that it has at least one fixed point on the tropical boundary $P(\mathcal{X}(\mathbb{R}^t) \setminus \mathcal{X}(\mathbb{R}^t)_+)$ from Proposition 2.13. It would be interesting to find an analogue of the *pA-pair* for a cluster-pA class which satisfies an appropriate condition, as we find in the surface theory (see Definition 3.10).

3. BASIC EXAMPLES: SL_2 -TEICHMÜLLER SEEDS

In this section we describe an important family of examples called the SL_2 -Teichmüller seeds, which are strongly related to the Teichmüller theory. A geometric description of the positive real parts and the tropical spaces associated with the SL_2 -Teichmüller seed is presented in Appendix B, which is used in Sections 3.2 and 3.3. In Section 3.2 we prove that the SL_2 -Teichmüller seeds are of Teichmüller type. In Section 3.3, we compare the Nielsen-Thurston types defined in Section 2 with the classification of mapping classes. In the case of the SL_2 -Teichmüller seed, the characterization of periodic classes described in Proposition 2.3 is complete. We show that cluster-reducible classes are reducible.

3.1. Definition of the seed. A *marked hyperbolic surface* is a pair (F, M) , where $F = F_{g,r}^s$ is an oriented surface of genus g with s punctures and r boundary components satisfying $2 - 2g - s < 0$ and $s + r > 0$, and $M \subset \partial F$ is a finite subset such that each boundary component has at least one point in M . The punctures together with elements of M are called *marked points*. We denote a marked hyperbolic surface by $F_{g,\vec{\delta}}^s$, where $\vec{\delta} = (\delta_1, \dots, \delta_r)$, $\delta_i := |M \cap \partial_i|$ indicates the number of marked points on the i -th boundary component. A connected component of $\partial F \setminus M$ is called a *boundary segment*. We denote the set of boundary segments by $B(F)$, and fix a numbering on its elements. Note that $|B(F)| = D$, where $D := \sum_{i=1}^r \delta_i$.

Definition 3.1 (the seed associated with an ideal triangulation).

- (1) An ideal arc on F is an isotopy class of an embedded arc connecting marked points, which is not isotopic to a puncture or an arc connecting two consecutive marked points on a common boundary component. An ideal triangulation of F is a family $\Delta = \{\alpha_i\}_{i=1}^n$ of ideal arcs, such that each connected component of $F \setminus \bigcup \alpha_i$ is a triangle whose vertices are marked points of F . One can verify that such an triangulation exists and that $n = 6g - 6 + 3r + 2s + D$ by considering the Euler characteristic.

- (2) For an ideal triangulation Δ of F , we define a skew-symmetric seed $\mathbf{i}_\Delta = (\Delta \cup B(F), B(F), \epsilon = \epsilon_\Delta)$ as follows. For an arc α of Δ which is contained in a self-folded triangle in Δ as in Figure 5, let $\pi_\Delta(\alpha)$ be the loop enclosing the triangle. Otherwise we set $\pi_\Delta(\alpha) := \alpha$. Then for a non-self-folded triangle τ in Δ , we define

$$\epsilon_{ij}^\tau := \begin{cases} 1, & \text{if } \tau \text{ contains } \pi_\Delta(\alpha_i) \text{ and } \pi_\Delta(\alpha_j) \text{ on its boundary in this clockwise order,} \\ -1, & \text{if the same holds, with the anti-clockwise order,} \\ 0, & \text{otherwise.} \end{cases}$$

Finally we define $\epsilon_{ij} := \sum_\tau \epsilon_{ij}^\tau$, where the sum runs over non-self-folded triangles in Δ .




FIGURE 5. Self-folded triangle

For an arc α of an ideal triangulation Δ which is a diagonal of an immersed quadrilateral in F (in this case the quadrilateral is unique), we get another ideal triangulation $\Delta' := (\Delta \setminus \{\alpha\}) \cup \{\beta\}$ by replacing α by the other diagonal β of the quadrilateral. We call this operation the *flip* along the arc α . One can directly check that the flip along the arc α_k corresponds to the mutation of the corresponding seed directed to the vertex k .

Theorem 3.2 (Whitehead (cf. [Pen12])). *Any two ideal triangulations of F are connected by a finite sequence of flips and relabelings.*

Hence the equivalence class of the seed \mathbf{i}_Δ is determined by the marked hyperbolic surface F , independent of the choice of the ideal triangulation. We refer to the seed \mathbf{i}_Δ as the SL_2 -Teichmüller seed on F . We denote the resulting cluster ensemble by $p = p_{SL_2, F} : \mathcal{A}_{SL_2, F} \rightarrow \mathcal{X}_{SL_2, F}$, the cluster modular group by $\Gamma_{SL_2, F} := \Gamma_{|\mathbf{i}_\Delta|}$, etc. The rank and the mutable rank of the SL_2 -Teichmüller seed are $N = |\Delta \cup B(F)| = 6g - 6 + 3r + 2s + 2D$ and $n = |\Delta| = 6g - 6 + 3r + 2s + D$, respectively.

Remark 3.3. Similarly one can define other seeds, which we shall call SL_m -Teichmüller seeds ($m \geq 2$), using an ideal triangulation as described in [FG06] Section 9.1. These are related to the moduli space of SL_m (or PGL_m)-local systems over F .

Though a flip induces a mutation, not every mutation is realized by a flip. Indeed, the existence of an arc contained in a self-folded triangle prevents us from performing the flip along such an arc. Therefore we generalize the concept of ideal triangulations, following [FST08].

Definition 3.4 (tagged triangulations).

- (1) A tagged arc on F is an ideal arc together with a label {plain, notched} assigned to each of its end, satisfying the following conditions:
 - the arc does not cut out a once-punctured monogon as in Figure 5,
 - each end which is incident to a marked point on the boundary is labelled plain, and
 - both ends of a loop are labelled in the same way.

The labels are called *tags*.

- (2) The *tagged arc complex* $\text{Arc}^{\bowtie}(F)$ is the clique complex for an appropriate compatibility relation on the set of tagged arcs on F . Namely, the vertices are tagged arcs and the collection $\{\alpha_1, \dots, \alpha_k\}$ spans a k -simplex if and only if they are mutually compatible. See, for the definition of the compatibility, [FST08]. The maximal simplices are called *tagged triangulations* and the codimension 1 simplices are called *tagged flips*.

Note that if the surface F has no punctures, then each tagged triangulation has only plain tags. If F has at least two punctures, then the tagged arc complex typically contains a cycle (which we call a \diamond -cycle) shown in the right of Figure 6. Here by convention, the plain tags are omitted in the diagram while the notched tags are represented by the \bowtie symbol. Compare with the ordinary arc complex, shown in the left of Figure 6. Compatibility relation implies that for a compatible set of tagged arcs and each puncture a , either one of the followings hold.

- (a) All tags at the puncture a are plain.
- (b) All tags at the puncture a are notched.
- (c) The number of arcs incident to the puncture a is at most two, and their tags at the puncture a is different.




FIGURE 6. \diamond -cycle

Definition 3.5 (the seed associated with a tagged triangulation). For a tagged triangulation Δ , let Δ° be an ideal triangulation obtained as follows:

- replace all tags at a puncture a of type (b) by plain ones, and
- for each puncture a of type (c), replace the arc α notched at a (if any) by a loop enclosing a and α .

A tagged triangulation Δ which satisfies $\delta_\Delta(a) = 1$ for all puncture a is naturally identified with the corresponding ideal triangulation Δ° . For a tagged triangulation Δ with a fixed numbering on the member arcs, we define a skew-symmetric seed by $\mathbf{i}_\Delta = (\Delta \cup B(F), B(F), \epsilon := \epsilon_{\Delta^\circ})$.

Then we get a complete description of the cluster complex associated with the SL_2 -Teichmüller seed in terms of ragged triangulations:

Theorem 3.6 (Fomin-Shapiro-Thurston [FST08] Proposition 7.10, Theorem 7.11).

For a marked hyperbolic surface $F = F_{g,\bar{\delta}}^s$, the tagged arc complex has exactly two

connected components (all plain/all notched) if $F = F_{g,0}^1$, and otherwise connected. The cluster complex of the SL_2 -Teichmüller seed on F is naturally identified with a connected component of the tagged arc complex $\text{Arc}^\bowtie(F)$ of the surface F . Namely,

$$\begin{cases} \mathcal{C}_{SL_2,F} \cong \text{Arc}(F) & \text{if } F = F_{g,0}^1, \\ \mathcal{C}_{SL_2,F} \cong \text{Arc}^\bowtie(F) & \text{otherwise.} \end{cases}$$

The coordinate groupoid of the SL_2 -Teichmüller seed is denoted by $\mathcal{M}^\bowtie(F)$, and called *tagged modular groupoid*. The subgroupoid $\mathcal{M}(F)$ whose objects are ideal triangulations and morphisms are (ordinary) flips is called the *modular groupoid*, which is described in [Pen12]. Next we see that the cluster modular group associated with the SL_2 -Teichmüller seed is identified with some extension of the mapping class group.

Definition 3.7 (the tagged mapping class group). The group $\{\pm 1\}^s$ acts on the tagged arc complex by alternating the tags at each puncture. The mapping class group naturally acts on the tagged arc complex, as well on the group $\{\pm 1\}^s$ by $(\phi_*\epsilon)(a) := \epsilon(\phi(a))$. Then the induced semidirect product $MC^\bowtie(F) := MC(F) \ltimes \{\pm 1\}^s$ is called the *tagged mapping class group*. The tagged mapping class group naturally acts on the tagged arc complex.

Proposition 3.8 (Bridgeland-Smith [BS15] Proposition 8.5 and 8.6). *The cluster modular group of the SL_2 -Teichmüller seed on F is naturally identified with the subgroup of the tagged mapping class group $MC^\bowtie(F)$ of F which consists of the elements that preserve connected components of $\text{Arc}^\bowtie(F)$. Namely,*

$$\begin{cases} \Gamma_{SL_2,F} \cong MC(F) & \text{if } F = F_{g,0}^1, \\ \Gamma_{SL_2,F} \cong MC^\bowtie(F) & \text{otherwise.} \end{cases}$$

We give a sketch of the construction of the isomorphism here, for later use.

sketch of the construction. Let us first consider the generic case $F \neq F_g^1$. Fixing a tagged triangulation Δ , we can think of the cluster modular group as $\Gamma_{SL_2,F} = \pi_1(\mathcal{M}^\bowtie(F), \Delta)$. For a mapping class $\psi = (\phi, \epsilon) \in MC^\bowtie(F)$, there exists a sequence of tagged flips $\mu_{i_1}, \dots, \mu_{i_k}$ from Δ to $\epsilon \cdot \phi^{-1}(\Delta)$ by Theorem 3.6. Since both ϕ and ϵ preserves the exchange matrix of the tagged triangulation, there exists a seed isomorphism $\sigma : \epsilon \cdot \phi^{-1}(\Delta) \rightarrow \Delta$. Then $I(\psi) := \sigma \circ \mu_{i_k} \cdots \mu_{i_1}$ defines an element of the cluster modular group. Hence we get a map $I : MC^\bowtie(F) \rightarrow \Gamma_{SL_2,F}$, which in turn gives an isomorphism. Since each element of $MC(F)$ preserves the tags, the case of $F = F_g^1$ is clear. \square

3.2. The SL_2 -Teichmüller seed on an marked hyperbolic surface is of Teichmüller type.

Theorem 3.9. *The SL_2 -Teichmüller seeds are of Teichmüller type.*

Proof. Condition (T1). We claim that the action of the cluster modular group on each positive space is properly discontinuous. Then the assertion follows from Proposition 2.7. First consider the action on the \mathcal{X} -space. By Proposition B.9, the action of the subgroup $MC(F) \subset \Gamma_{SL_2,F}$ on the \mathcal{X} -space $\mathcal{X}(\mathbb{R}_{>0})$ coincide with the geometric action. Hence this action of $MC(F)$ is properly discontinuous, as is well-known. See, for instance, [FM12]. From the definition of the action of $\Gamma_{SL_2,F} = MC^\bowtie(F)$ on the tagged arc complex and eq. (1.1), one can verify that an element $(\phi, \epsilon) \in MC^\bowtie(F)$ acts on the positive \mathcal{X} -space as $(\phi, \epsilon)g = \phi(\iota(\epsilon)g)$, where $\iota(\epsilon) := \prod_{\epsilon(p)=-1} \iota_p$ is a composition of the involutions defined in Definition B.7. Now suppose that there exists a compact set $K \subset \mathcal{X}(\mathbb{R}_{>0})$ and an infinite sequence $\psi_n = (\phi_n, \epsilon_n) \in \Gamma_{SL_2,F}$ such that $\psi_n(K) \cap K \neq \emptyset$. Since $\{\pm 1\}^s$ is a finite group,

$(\phi_n) \subset MC(F)$ is an infinite sequence and there exists an element $\epsilon \in \{\pm 1\}^s$ such that $\epsilon_n = \epsilon$ for infinitely many n . Hence we have

$$\emptyset \neq \psi_n(K) \cap K = \phi_n(\iota(\epsilon)K) \cap K \subset \phi_n(\iota(\epsilon)K \cup K) \cap (\iota(\epsilon)K \cup K)$$

for infinitely many n , which is a contradiction to the proper discontinuity of the action of $MC(F)$. Hence the action of $\Gamma_{SL_2, F}$ on the \mathcal{X} -space is properly discontinuous. The action on the \mathcal{A} -space is similarly shown to be properly discontinuous. Here the action of ϵ is described as $\iota'(\epsilon) := \prod_{\epsilon(p)=-1} \iota'_p$, where ι'_p is the involution changing the horocycle assigned to the puncture p to the conjugated one (see [FT12]).

Condition (T2). Note that for a tagged triangulation $\Delta = \{\gamma_1, \dots, \gamma_N\}$ without digons as in the left of Figure 7 in Appendix B, the map $\Psi_\Delta|_{[S_\Delta]}$ is given by $\Psi([w_1, \dots, w_N]) = (\sqcup w_j \gamma_j, \pm)$, where the sign at a puncture p is defined to be $+1$ if the tags of arcs at p are plain, and -1 if the tags are notched. Then on the image of these maps, the equivalence condition holds. Let us consider the tagged triangulation Δ_j in the \diamond -cycle, see Figure 6. From the definition of the tropical \mathcal{X} -transformations, we have

$$\begin{cases} x_{\Delta_2}(\alpha) = -x_{\Delta_1}(\alpha) \\ x_{\Delta_2}(\beta) = x_{\Delta_1}(\beta) \end{cases} \quad \begin{cases} x_{\Delta_3}(\alpha) = -x_{\Delta_1}(\alpha) \\ x_{\Delta_3}(\beta) = -x_{\Delta_1}(\beta) \end{cases} \quad \begin{cases} x_{\Delta_4}(\alpha) = x_{\Delta_1}(\alpha) \\ x_{\Delta_4}(\beta) = -x_{\Delta_1}(\beta) \end{cases}.$$

Hence the equivalence condition on the image of the \diamond -cycle holds. \square

3.3. Comparison with the Nielsen-Thurston classification of elements of the mapping class group. Let F be a hyperbolic surface of type F_g^1 or $F_{g,\delta}$ throughout this subsection. Recall that in this case we have $\Gamma_{SL_2, F} \cong MC(F)$ and $\mathcal{C}_{SL_2, F} \cong \text{Arc}(F)$, see Proposition 3.8 and Theorem 3.6. Let us recall the Nielsen-Thurston classification.

Definition 3.10 (Nielsen-Thurston classification). A mapping class $\phi \in MC(F)$ is said to be

- (1) *reducible* if it fixes an isotopy class of a finite union of mutually disjoint simple closed curves on F , and
- (2) *pseudo-Anosov* (pA) if there is a pair of mutually transverse filling laminations $G_\pm \in \mathcal{ML}_0^+(F)$ and a scalar factor $\lambda > 0$ such that $\phi(G_\pm) = \lambda^{\pm 1} G_\pm$. The pair of projective laminations $[G_\pm]$ is called the pA -pair of ϕ .

Here a lamination $G \in \widetilde{\mathcal{ML}}(F)$ is said to be *filling* if each component of $F \setminus G$ is unpunctured or once-punctured polygon. It is known (see, for instance, [FLP12]) that each mapping class is at least one of periodic, reducible, or pA , and a pA class is neither periodic nor reducible. Furthermore a mapping class ϕ is reducible if and only if it fixes a non-filling projective lamination, and is pA if and only if it satisfies $\phi(G) = \lambda G$ for some filling lamination $G \in \mathcal{ML}_0^+(F)$ and a scalar $\lambda > 0, \neq 1$. A pA class ϕ has the following asymptotic behavior of orbits in $\mathcal{PML}_0(F)$: for any projective lamination $[G] \in \mathcal{PML}_0(F)$ we have $\lim_{n \rightarrow \infty} \phi^{\pm n}[G] = [G_\pm]$.

We shall start with periodic classes. In this case the characterization of periodic classes described in Proposition 2.3 is complete:

Proposition 3.11. *For a mapping class $\phi \in \Gamma_{SL_2, F}$, the following conditions are equivalent.*

- (i) *The mapping class ϕ fixes a cell $C \in \mathcal{C}$ of finite type.*
- (ii) *The mapping class ϕ is periodic.*
- (iii) *The mapping class ϕ has fixed points in $\mathcal{A}_{SL_2, F}(\mathbb{R}_{>0})$ and $\mathcal{X}_{SL_2, F}(\mathbb{R}_{>0})$.*

Lemma 3.12. *The cells of finite type (see Proposition 2.3) in the cluster complex are in one-to-one correspondence with ideal cell decompositions of F . Here an*

ideal cell decomposition is a family $\Delta = \{\alpha_i\}$ of ideal arcs such that each connected component of $F \setminus \bigcup \alpha_i$ is a polygon.

Proof. Let $C = (\alpha_1, \dots, \alpha_k)$ be a cell in the cluster complex, which is represented by a family of ideal arcs. Suppose that $\{\alpha_1, \dots, \alpha_k\}$ is an ideal cell decomposition. Then supercells of C are obtained by adding some ideal arcs on the surface $F \setminus \bigcup_{i=1}^k \alpha_i$ to $\{\alpha_1, \dots, \alpha_k\}$, which are finite since such an ideal arc must be a diagonal of a polygon. Conversely suppose that $\{\alpha_1, \dots, \alpha_k\}$ is not an ideal cell decomposition. Then there exists a connected component F_0 of $F \setminus \bigcup_{i=1}^k \alpha_i$ which has a half-twist or a Dehn twist in its mapping class group. Hence F_0 has infinitely many ideal triangulations, consequently C has infinitely many supercells. \square

proof of Proposition 3.11. It suffices to show that the condition (iii) implies the condition (i). Let \mathcal{C}^* denote the union of all cells of finite type in the cluster complex. In view of Lemma 3.12, Penner's convex hull construction ([Pen12]Chapter 4) gives a mapping class group equivariant isomorphism

$$\mathcal{C}^* \cong \widetilde{T}(F)/\mathbb{R}_{>0},$$

from which the assertion follows. \square

Next we focus on cluster-reducible classes and their relation with *reducible classes*. Observe that by Theorem 3.9 a mapping class is cluster-reducible if and only if it fixes an isotopy class of a finite union of mutually disjoint ideal arcs on F .

Proposition 3.13. *We have the followings.*

- (1) *A mapping class ϕ is cluster-reducible if and only if it fixes an unbounded lamination with real weights $L = (\bigsqcup w_j \gamma_j, \pm)$, where $w_j \in \mathbb{R}$. If ϕ is proper reducible, then it induces a mapping class on the surface obtained by cutting F along the multiarc $\bigsqcup \gamma_j$.*
- (2) *A cluster-reducible class is reducible.*
- (3) *A filling lamination is cluster-filling.*

Proof.

(1). The assertion follows from Proposition 2.13(2). Note that an element of $P\mathcal{X}(\mathbb{R}^t)_+$ consists of elements of the form $L = (\bigsqcup w_j \gamma_j, \pm)$, where $w_j \in \mathbb{R}_{>0}$.

(2). Let $\phi \in MC(F)$ be a cluster-reducible class, $L = (\bigsqcup w_j \gamma_j, \pm)$ a fixed lamination, and $\bigsqcup \gamma_j$ the corresponding multiarc. One can pick representatives of ϕ and γ so that $\phi(\gamma) = \gamma$ on F . Then by cutting F along $\bigsqcup \gamma_j$, we obtain a surface F' with boundary. Since ϕ fixes $\bigsqcup \gamma_j$, it induces a mapping class ϕ' on F' which may permute the boundary components. Let C' be the multicurve isotopic to the boundary of F' . Since ϕ' fixes C' , the preimage C of C' in F is fixed by ϕ . Therefore ϕ is reducible.

(3) Let G be a non-cluster-filling lamination. Let γ be an ideal arc such that $a_\gamma(G) = 0$. Then G has no intersection with γ . Since G has compact support, there is a twice-punctured disk which surrounds γ and disjoint from G , which implies that G is non-filling. \square

Example 3.14 (a reducible class which is not cluster-reducible). Let C be a non-separating simple closed curve in $F = F_g^s$, and $\phi \in MC(F)$ a mapping class given by the Dehn twist along C on a tubular neighborhood $\mathcal{N}(C)$ of C and a pA class on $F \setminus \mathcal{N}(C)$. Then ϕ is a reducible class which is not cluster-reducible.

Proof. The reducibility is clear from the definition. Let $F' := F \setminus \mathcal{N}(C)$. If ϕ fixes an ideal arc contained in F' , then by Proposition 3.13 we see that the restriction $\phi|_{F'} \in MC(F')$ is reducible, which is a contradiction. Moreover since ϕ is the Dehn

twist along C near the curve C , it cannot fix ideal arcs which traverse the curve C . Hence ϕ is cluster-irreducible. \square

Example 3.15 (a cluster-filling lamination which is not filling). Let C be a simple closed curve in $F = F_g^s$, and $\{P_j\}$ be a pants decomposition of F which contains C as a decomposing curve: $F = \bigcup_j P_j$. For a component P_j which contains a puncture, let $G_j \in \mathcal{ML}_0^+(P_j)$ be a filling lamination such that $i(G_j, C) = 0$. For a component P_j which does not contain any punctures, choose an arbitrary lamination $G_j \in \mathcal{ML}_0^+(P_j)$. Then $G := \bigsqcup_j G_j \sqcup C \in \mathcal{ML}_0^+(F)$ is a cluster-filling lamination which is not filling.

Just as the fact that a mapping class is reducible if and only if it fixes a non-filling projective lamination, we expect that a mapping class is cluster-reducible if and only if it fixes a non-cluster-filling projective lamination.

APPENDIX A. THE SEED OF TYPE L_k IS OF TEICHMÜLLER TYPE

Here we show that the seed of type L_k is of Teichmüller type. Recall that $\Gamma_{L_k} \cong \mathbb{Z}$ from Example 2.8 and the generator ϕ is a cluster Dehn twist.

Theorem A.1. *The seed of type L_k is of Teichmüller type.*

Proof. Condition (T1). Since the cluster complex \mathcal{C}_{L_k} is homeomorphic to the real line and the cluster modular group acts by the shift, it suffices to show that each orbit of the generator ϕ is divergent. In the case of the \mathcal{X} -space, let us consider the following recurrence relation:

$$\begin{aligned} X_0^{(n)} &= X_1^{(n-1)}(1 + X_0^{(n-1)})^k \\ X_1^{(n)} &= (X_0^{(n-1)})^{-1}. \end{aligned}$$

We claim that $\log X_0^{(n)}$ and $\log X_1^{(n)}$ diverges as $n \rightarrow \infty$. Setting $x_n := \log X_0^{(n)}$ and deleting $X_1^{(n)}$, we have a 3-term recurrence relation

$$x_n = -x_{n-2} + k \log(1 + \exp x_{n-1}).$$

Subtracting x_{n-1} from the both sides, we have

$$(A.1) \quad y_n = y_{n-1} + f(x_{n-1})$$

where we set $y_n := x_n - x_{n-1}$ and $f(x) := k \log(1 + \exp x) - 2x$. Since $f(x)$ is positive, if y_N is non-negative for some N , then $(y_n)_{n \geq N}$ is monotone increasing and

$$x_n = x_{N-1} + \sum_{k=N}^n y_k \geq x_{N-1} + (n - N + 1)y_1 \rightarrow +\infty$$

as $n \rightarrow \infty$. Therefore we claim that y_N is non-negative for some N .

Suppose that $y_n < 0$ for all $n \geq 1$. Note that if $x_n \leq 0$ for some n , then $x_{n+2} > 0 > x_{n+1}$, hence $y_{n+2} > 0$. Therefore it suffices to consider the case $x_n > 0$ for all $n \geq 1$. Then x_n is a decreasing sequence of positive numbers. In this case, from eq. (A.1) we have

$$y_n = y_1 + \sum_{k=1}^{n-1} f(x_k) \geq y_1 + (n-1) \min_{0 \leq x \leq x_1} f(x) \rightarrow \infty$$

as $n \rightarrow \infty$, which is a contradiction. Thus $\log X_0^{(n)}$ diverges to $+\infty$ and $\log X_1^{(n)}$ diverges to $-\infty$. Hence the condition (T1) holds for the \mathcal{X} -space. We have proved the case of the \mathcal{A} -space in Theorem 2.33.

Condition (T2). In the tropical \mathcal{X} -coordinate (x_0, x_1) associated with the seed \mathbf{i}_k , the action of ϕ on the tropical \mathcal{X} -space is expressed as follows:

$$\phi(x_0, x_1) = (x_1 + k \max\{0, x_0\}, -x_0).$$

To prove the condition (T2), we need to know the change of signs of tropical coordinates induced by the action described above. The following lemma follows from a direct calculation.

Lemma A.2. *Consider the following recurrence relation:*

$$\begin{cases} x_0^{(n)} &= x_1^{(n-1)} + kx_0^{(n-1)}, \\ x_1^{(n)} &= -x_0^{(n-1)}. \end{cases}$$

Then we have that if $x_0^{(0)} > 0$ and $x_0^{(0)} + x_1^{(0)} > 0$, then $x_0^{(n)} > 0$ and $x_0^{(n)} + x_1^{(n)} > 0$ for all $n \geq 0$.

In particular, the tropical action of ϕ on the cone $C_+ := \{(x_0, x_1) \mid x_0 > 0, x_0 + x_1 > 0\}$ is expressed by the linear transformation

$$(A.2) \quad \phi \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \begin{pmatrix} k & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix}.$$

Let $L \in \mathcal{X}(\mathbb{R}^t)_+ \setminus \{0\}$ be an arbitrary point, and \mathbf{i} a non-negative seed for L . Then $L = (x_0, x_1)$ and $x_0, x_1 \geq 0$ in the coordinate associated with \mathbf{i} . In particular we have $x_0 + x_1 > 0$.

If $x_0 > 0$, by Lemma A.2 we have $x_1^{(n)} = -x_0^{(n-1)} < 0$ for all $n \geq 1$, which implies that no seed other than \mathbf{i} is non-negative for L .

If $x_0 = 0$, we have $x_1 > 0$ and $x_0^{(1)} > 0$, which implies that $\mu_0(\mathbf{i})$ is again a non-negative seed for L , while any other seeds are not non-negative for L from the argument in the previous paragraph. Hence the condition (T2) holds. \square

Next we study the asymptotic behavior of orbits of the generator ϕ of Γ_{L_k} on the tropical \mathcal{X} -space, which may be related with that of general cluster Dehn twists.

Proposition A.3. *For $k \geq 2$, the generator ϕ of the cluster modular group Γ_{L_k} has unique attracting/repelling fixed points $[L_{\pm}] \in P\mathcal{X}(\mathbb{R}^t)$ such that for all $L \in \mathcal{X}(\mathbb{R}^t)$ we have*

$$\lim_{n \rightarrow \infty} \phi^{\pm n}([L]) = [L_{\pm}] \text{ in } P\mathcal{X}(\mathbb{R}^t).$$

Proof. Note that $\phi^{-1}(x_0, x_1) = (-x_1, x_0 + k \min\{0, x_1\})$. By a similar argument as Lemma A.2, we have that the cone $C_- := \{(x_0, x_1) \mid x_1 < 0, x_0 + x_1 < 0\}$ is stable under ϕ^{-1} and on this cone

$$(A.3) \quad \phi^{-1} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & k \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix}.$$

Together with the fact that $\phi(-1, 0) = (0, 1)$ and $\phi(0, 1) = (1, 0)$, we see that for all $L \in \mathcal{X}(\mathbb{R}^t)$, $\phi^{\pm N}(L) \in C_{\pm}$ for a sufficiently large number N . Then each orbit $(\phi^n(L))_{n \geq 0}$ projectively converges to the unique attracting fixed point $[L_+]$ of the linear action (A.2), which is represented by $(k + \sqrt{k^2 - 4}, -2)$. Similarly $(\phi^n(L))_{n \leq 0}$ projectively converges to the unique repelling fixed point $[L_-]$ of the linear action (A.3), which is represented by $(k - \sqrt{k^2 - 4}, -2)$. \square

APPENDIX B. THE POSITIVE REAL PARTS AND THE TROPICAL SPACES ASSOCIATED WITH THE SL_2 -TEICHMÜLLER SEED

Here we give a geometric description of the positive real parts and the tropical spaces associated with the SL_2 -Teichmüller seed on a marked hyperbolic surface $F = F_{g,\delta}^s$. Most of the contents of this section seems to be well-known to specialists,

but they are scattered in literature. Therefore we tried to gather the result and give a coherent presentation of the data associated with SL_2 -Teichmüller seeds.

B.1. Positive spaces and the Teichmüller spaces. Here we describe the positive real parts of $\mathcal{A}_{SL_2, F}$ and $\mathcal{X}_{SL_2, F}$ geometrically. Main references are [FG07, FT12, Pen12]. For simplicity, we only deal with the case of empty boundary, $r = 0$. The case of non-empty boundary is reduced to the case of empty boundary by duplicating the surface and considering the invariant subspace of the Teichmüller/lamination spaces under the natural involution. See, for details, [Pen12] section 2.

Let $F = F_g^s$ be a hyperbolic punctured surface. Let $T(F)$ denote the Teichmüller space of all complete finite-area hyperbolic structures on F . Namely,

$$T(F) := \text{Hom}'(\pi_1(F), \text{PSL}_2(\mathbb{R}))/\text{PSL}_2(\mathbb{R}),$$

where $\text{Hom}'(\pi_1(F), \text{PSL}_2(\mathbb{R}))$ consists of faithful representations $\rho : \pi_1(F) \rightarrow \text{PSL}_2(\mathbb{R})$ such that

- (1) the image of ρ is a discrete subgroup of $\text{PSL}_2(\mathbb{R})$, and
- (2) it maps every peripheral loops to parabolic elements, essential ones to hyperbolic ones.

Definition B.1 (decorated Teichmüller space). The trivial bundle $\tilde{T}(F) := T(F) \times \mathbb{R}_{>0}^s$ is called the *decorated Teichmüller space*. Let $\varpi : \tilde{T}(F) \rightarrow T(F)$ be the natural projection.

Proposition B.2 (Penner [Pen12] Chapter 2, Theorem 2.5). *For any ideal triangulation Δ of F , there exists a real analytic diffeomorphism*

$$A_\Delta : \tilde{T}(F) \rightarrow \mathbb{R}_{>0}^\Delta$$

which we call the Penner coordinates with respect to Δ . Furthermore the Penner coordinates give rise to a positive space $\psi_\Delta : \mathcal{M}(F) \rightarrow \text{Pos}(\mathbb{R})$.

The Penner coordinate is given by assigning the *lambda length* to each ideal arc. In [FT12], the authors generalized the definition of the \mathcal{A} -coordinates to tagged arcs:

Theorem B.3 (Fomin-Thurston [FT12] Theorem 8.6). *The above functor extends to a positive space $\psi_\Delta^\sqcap : \mathcal{M}^\sqcap(F) \rightarrow \text{Pos}(\mathbb{R})$ and the positive real part is naturally identified with the decorated Teichmüller space $\tilde{T}(F)$, i.e., $\mathcal{A}(\mathbb{R}_{>0}) \cong \tilde{T}(F)$.*

Definition B.4 (the enhanced Teichmüller space). Let $T(F)'$ denote the Teichmüller space of all complete (not necessarily finite-area) hyperbolic structures on F . Namely,

$$T(F)' := \text{Hom}''(\pi_1(F), \text{PSL}_2(\mathbb{R}))/\text{PSL}_2(\mathbb{R}),$$

where $\text{Hom}''(\pi_1(F), \text{PSL}_2(\mathbb{R}))$ consists of faithful representations $\rho : \pi_1(F) \rightarrow \text{PSL}_2(\mathbb{R})$ such that

- (1) the image of ρ is a discrete subgroup of $\text{PSL}_2(\mathbb{R})$, and
- (2) it maps every peripheral loops to parabolic or hyperbolic elements, essential ones to hyperbolic ones.

The *enhanced Teichmüller space* $\widehat{T}(F)$ is defined to be the 2^s -folded branched cover of $T(F)'$, whose fiber over a point $\rho \in T(F)'$ is the collection of all tuples of orientations on the punctures which are represented hyperbolically under ρ .

Note that for a point of $T(F)$, each puncture is represented parabolically. Hence there is a natural embedding $\iota : T(F) \rightarrow \widehat{T}(F)$.

Proposition B.5 (Fock-Goncharov [FG07] Section 4.1). *For any ideal triangulation Δ of F , there exists a real analytic diffeomorphism*

$$X_\Delta : \widehat{T}(F) \rightarrow \mathbb{R}_{>0}^\Delta$$

which we call the Fock-Goncharov coordinates with respect to Δ . Furthermore the Fock-Goncharov coordinates give rise to a positive space $\psi_\chi : \mathcal{M}(F) \rightarrow \text{Pos}(\mathbb{R})$.

Proposition B.6. *The above functor extends to a positive space $\psi_\chi^\Delta : \mathcal{M}^\Delta(F) \rightarrow \text{Pos}(\mathbb{R})$ and the positive real part is naturally identified with the enhanced Teichmüller space $\widehat{T}(F)$, i.e., $\mathcal{X}(\mathbb{R}_{>0}) \cong \widehat{T}(F)$.*

Although the above proposition seems to be well-known to specialists, we could not find any proof in literature. Therefor we give a proof here for completeness.

Definition B.7 (the coordinates associated with a tagged triangulation). We already have the Fock-Goncharov coordinates $X_\Delta : \mathcal{X}(\mathbb{R}_{>0}) \rightarrow \mathbb{R}_{>0}^\Delta$ with respect to any ideal triangulation Δ . We define a coordinate system for any tagged triangulations by the following conditions:

- (1) Suppose two tagged triangulations Δ_1, Δ_2 coincide except for the tags at a puncture p . Then we set

$$X_{\Delta_1}(g, \alpha_1) = X_{\Delta_2}(\iota_p(g), \alpha_2)$$

for all $g \in \mathcal{X}(\mathbb{R}_{>0})$, where $\alpha_i \in \Delta_i$ ($i = 1, 2$) are the corresponding arcs, ι_p is the involution on $\mathcal{X}(\mathbb{R}_{>0})$ reversing the fiber parameter of the cover assigned to the puncture p .

- (2) If the tags of a tagged triangulation Δ are all plain, then we set $X_\Delta(g, \alpha) := X_{\Delta^\circ}(g, \alpha^\circ)$ for all $g \in \mathcal{X}(\mathbb{R}_{>0})$ and $\alpha \in \Delta$. The right-hand side is the Fock-Goncharov coordinate for the ideal triangulation Δ° .
- (3) If a tagged triangulation Δ have a punctured digon shown in the left of Figure 7, then we set $X_\Delta(g, \gamma) := X_{\Delta^\circ}(g, \gamma^\circ)$ for $\gamma \neq \alpha$, and respecting the rule (1) we set

$$X_\Delta(g, \alpha) = X_{\Delta'}(\iota_p(g), \alpha') = X_{\Delta^\circ}(\iota_p(g), \beta^\circ).$$




FIGURE 7

The following is the key lemma to ensure that the above definition is well-defined, which is essentially a special case of Lemma 12.3 in [FG06]:

Lemma B.8 (Fock-Goncharov [FG06]). *In the notation of Figure 6, we have*

$$X_{\Delta_4}(g, \alpha)X_{\Delta_4}(\iota_p(g), \beta) = 1$$

for all $g \in \widehat{T}(F)$.

proof of Proposition B.6. We need to show that each coordinate transformations $X_{\Delta_i} \circ X_{\Delta_j}^{-1}$ in the \diamond -cycle coincide with the corresponding cluster \mathcal{X} -transformations.

$\Delta_1 \xleftrightarrow{\mu_\beta} \Delta_4$. Note that the coordinate transformations of X_γ ($\gamma \neq \alpha$) coincide with the cluster transformations, since they are Fock-Goncharov coordinates. In particular we have $X_{\Delta_1}(g, \beta) = X_{\Delta_4}(g, \beta)^{-1}$. Hence by Lemma B.8 we have

$$X_{\Delta_1}(g, \alpha) = X_{\Delta_1^\circ}(\iota_p(g), \beta^\circ) = X_{\Delta_4^\circ}(\iota_p(g), \beta^\circ)^{-1} = X_{\Delta_4^\circ}(g, \alpha^\circ) = X_{\Delta_4}(g, \alpha).$$

$\Delta_1 \xleftrightarrow{\mu_\alpha} \Delta_2$. Note that the coordinate transformations of X_γ ($\gamma \neq \beta$) coincide with the corresponding cluster transformations, and we have

$$X_{\Delta_2}(g, \beta) = X_{\Delta_4}(\iota_p(g), \alpha) = X_{\Delta_1}(\iota_p(g), \alpha) = X_{\Delta_1^\circ}(g, \beta^\circ) = X_{\Delta_1}(g, \beta).$$

The remaining cases follow from a symmetric argument. \square

The monomial morphism between the positive spaces of the SL_2 -Teichmüller seed coincides with $p = \iota \circ \varpi : \tilde{T}(F) \rightarrow \hat{T}(F)$ (see, for instance, [Pen12] Chapter 1, Corollary 4.16(c)). In particular, the \mathcal{U} -space is naturally identified with the Teichmüller space, *i.e.*, $\mathcal{U}(\mathbb{R}_{>0}) \cong T(F)$.

The mapping class group naturally acts on the Teichmüller space $T(F)$ and $T(F)'$ via the Dehn-Nielsen embedding [FM12] $MC(F) \rightarrow \text{Out}(\pi_1(F))$. These two actions extend to $\tilde{T}(F)$ and $\hat{T}(F)$ by permuting the fiber parameters according to the action on the punctures.

Proposition B.9 (Penner [Pen12] Chapter 2, Theorem 2.10). *For any ideal triangulation Δ of F and a mapping class $\phi \in MC(F)$, the following diagrams commute :*

$$\begin{array}{ccc} \tilde{T}(F) & \xrightarrow{A_\Delta} & \mathbb{R}_{>0}^\Delta \\ \phi \downarrow & & \downarrow I(\phi) \\ \tilde{T}(F) & \xrightarrow{A_\Delta} & \mathbb{R}_{>0}^\Delta \end{array} \quad \begin{array}{ccc} \hat{T}(F) & \xrightarrow{X_\Delta} & \mathbb{R}_{>0}^\Delta \\ \phi \downarrow & & \downarrow I(\phi) \\ \hat{T}(F) & \xrightarrow{X_\Delta} & \mathbb{R}_{>0}^\Delta \end{array}$$

In particular, these natural actions coincide with the action as a subgroup of the cluster modular group of the SL_2 -Teichmüller seed. Compare with eq. (1.1).

B.2. Tropical spaces and the lamination spaces. Next we describe the tropical spaces geometrically, following [FG07]. As before, we focus on the case of empty boundary $r = 0$.

Definition B.10. A *decorated rational (bounded) lamination* on F is an isotopy class of a disjoint union of simple closed curves in F with rational numbers (called weights) assigned to each curve so that the weight is positive unless the corresponding curve is peripheral. Each curve is called a *leaf* of the lamination.

We denote a decorated rational lamination by $L = \bigsqcup w_j \gamma_j$, and denote the set of decorated rational laminations by $\tilde{\mathcal{L}}(F; \mathbb{Q})$. We refer to the weighted peripheral leaves as *decorations* on the lamination.

We call a decorated rational lamination with no decoration (*i.e.* every weights of peripheral curves are 0) as a *rational (bounded) lamination*. Denote the set of rational laminations by $\mathcal{L}(F; \mathbb{Q})$. There is a canonical projection $\varpi : \tilde{\mathcal{L}}(F; \mathbb{Q}) \rightarrow \mathcal{L}(F; \mathbb{Q})$ forgetting decorations.

Following [FG07], we associate a rational number with an ideal arc α . For a decorated rational lamination $L = \bigsqcup w_j \gamma_j$, isotope each curve γ_j so that the intersection with α is minimal. Then define $a_\alpha(L) := \sum_j w_j i(\gamma_j, \alpha)$, where i denotes the geometric intersection number.

Proposition B.11 (Fock-Goncharov [FG07] Section 3.2). *For any ideal triangulation Δ of F , the map*

$$a_\Delta : \tilde{\mathcal{L}}(F; \mathbb{Q}) \rightarrow \mathbb{Q}^\Delta; L \mapsto \{a_\alpha(L)\}_{\alpha \in \Delta}$$

gives a bijection.

For a flip along $\alpha \in \Delta$, the corresponding change of the above coordinates coincide with the tropical cluster \mathcal{A} -transformation. Since the tropical cluster \mathcal{A} -transformation is continuous with respect to the standard topology on \mathbb{Q}^N , the coordinates induce a topology on $\tilde{\mathcal{L}}(F; \mathbb{Q})$.

Definition B.12 (real lamination spaces). Define the *real decorated lamination space* $\tilde{\mathcal{L}}(F; \mathbb{R})$ as the completion of $\tilde{\mathcal{L}}(F; \mathbb{Q})$ with respect to the topology induced by the coordinates defined above. Similarly define $\mathcal{L}(F; \mathbb{R})$ as the completion of $\mathcal{L}(F; \mathbb{Q})$.

The coordinates defined above give rise to a functor $\mathcal{M}(F) \rightarrow \text{PL}$.

Proposition B.13 (Fomin-Thurston [FT12]). *The coordinate functor defined above naturally extends to the tagged modular groupoid $\mathcal{M}^\square(F)$, and the tropical space of the positive space $\psi_A^\square : \mathcal{M}^\square(F) \rightarrow \text{Pos}(\mathbb{R})$ given in Theorem B.3 is naturally identified with the real decorated lamination space $\tilde{\mathcal{L}}(F; \mathbb{R})$, i.e., $\mathcal{A}(\mathbb{R}^t) \cong \tilde{\mathcal{L}}(F; \mathbb{R})$.*

Although the geometric meaning of irrational points in $\tilde{\mathcal{L}}(F; \mathbb{R})$ is not so clear from the above definition, we have the following result.

Theorem B.14 (for instance, [PH92]). *There are natural PL homeomorphisms*

$$\begin{aligned} \mathcal{L}(F; \mathbb{R}) &\cong \mathcal{ML}_0^+(F), \\ \tilde{\mathcal{L}}(F; \mathbb{R}) &\cong \widetilde{\mathcal{ML}}(F) \end{aligned}$$

where $\mathcal{ML}_0^+(F) := \mathcal{ML}_0(F) \cup \{\emptyset\}$ is the space of measured geodesic laminations with compact supports attached with the empty lamination, and $\widetilde{\mathcal{ML}}(F) := \mathcal{ML}_0^+(F) \times \mathbb{R}^s$ is a trivial bundle. See, for details, [PH92] [PP93]. Moreover, the bundle projection $\widetilde{\mathcal{ML}}(F) \rightarrow \mathcal{ML}_0^+(F)$ coincides with the projection $\varpi : \tilde{\mathcal{L}}(F; \mathbb{R}) \rightarrow \mathcal{L}(F; \mathbb{R})$.

Definition B.15. A *rational unbounded lamination* consists of the following data:

- (1) an isotopy class of a disjoint union of simple closed curves and ideal arcs $\{\gamma_j\}_j$ in F with positive rational weights $\{w_j\}$ assigned to each curve.
- (2) a tuple of orientations on each puncture to which some curves incident.

We denote these data by $L = (\bigsqcup w_j \gamma_j, \pm)$.

Denote the set of rational unbounded laminations by $\hat{\mathcal{L}}(F; \mathbb{Q})$. We have a natural embedding $\iota : \mathcal{L}(F; \mathbb{Q}) \hookrightarrow \hat{\mathcal{L}}(F; \mathbb{Q})$, where the orientation data is unnecessary since the leaves of a bounded lamination do not incident to any punctures.

Proposition B.16 (Fock-Goncharov [FG07] Section 3.1). *For any ideal triangulation Δ of F , there exists a natural bijection*

$$x_\Delta : \hat{\mathcal{L}}(F; \mathbb{Q}) \rightarrow \mathbb{Q}^\Delta.$$

For a flip along $\alpha \in \Delta$, the corresponding change of the above coordinates coincide with the tropical cluster \mathcal{X} -transformation. By the continuity of the tropical cluster \mathcal{X} -transformations, we can define the real unbounded lamination space $\hat{\mathcal{L}}(F; \mathbb{R})$ as the completion of $\hat{\mathcal{L}}(F; \mathbb{Q})$. Hence we get a functor $\mathcal{M}(F) \rightarrow \text{PL}$.

Proposition B.17 (Fomin-Thurston [FT12] Theorem 13.6). *The coordinate functor defined above naturally extends to the tagged modular groupoid $\mathcal{M}^{\bowtie}(F)$, and the tropical space of the positive space $\psi_{\mathcal{X}} : \mathcal{M}(F) \rightarrow \text{Pos}(\mathbb{R})$ given in Proposition B.2 is naturally identified with the real decorated lamination space $\widehat{\mathcal{L}}(F; \mathbb{R})$, i.e., $\mathcal{X}(\mathbb{R}^t) \cong \widehat{\mathcal{L}}(F; \mathbb{R})$.*

The extension is in the same manner as the one described in Proposition B.6.

Note that the mapping class group naturally acts on the lamination spaces. These natural actions coincide with the action as a subgroup of the cluster modular group, as in Proposition B.9.

REFERENCES

- [Bro82] Kenneth S. Brown, *Complete Euler characteristics and fixed-point theory*, J. Pure Appl. Algebra **24** (1982), no. 2, 103–121. MR 651839
- [BS15] Tom Bridgeland and Ivan Smith, *Quadratic differentials as stability conditions*, Publ. Math. Inst. Hautes Études Sci. **121** (2015), 155–278. MR 3349833
- [FG06] Vladimir V. Fock and Alexander B. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. (2006), no. 103, 1–211. MR 2233852
- [FG07] ———, *Dual Teichmüller and lamination spaces*, Handbook of Teichmüller theory. Vol. I, IRMA Lect. Math. Theor. Phys., vol. 11, Eur. Math. Soc., Zürich, 2007, pp. 647–684. MR 2349682
- [FG09] ———, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 6, 865–930. MR 2567745
- [FG16] ———, *Cluster Poisson varieties at infinity*, Selecta Math. (N.S.) **22** (2016), no. 4, 2569–2589. MR 3573965
- [FLP12] Albert Fathi, François Laudenbach, and Valentin Poénaru, *Thurston’s work on surfaces*, Mathematical Notes, vol. 48, Princeton University Press, Princeton, NJ, 2012, Translated from the 1979 French original by Djun M. Kim and Dan Margalit. MR 3053012
- [FM12] Benson Farb and Dan Margalit, *A primer on mapping class groups*, Princeton Mathematical Series, vol. 49, Princeton University Press, Princeton, NJ, 2012. MR 2850125
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston, *Cluster algebras and triangulated surfaces. I. Cluster complexes*, Acta Math. **201** (2008), no. 1, 83–146. MR 2448067
- [FST12] Anna Felikson, Michael Shapiro, and Pavel Tumarkin, *Skew-symmetric cluster algebras of finite mutation type*, J. Eur. Math. Soc. (JEMS) **14** (2012), no. 4, 1135–1180. MR 2928847
- [FT12] Sergey Fomin and Dylan Thurston, *Cluster algebras and triangulated surfaces. II. Lambda lengths*, preprint (2012).
- [FZ03] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. II. Finite type classification*, Invent. Math. **154** (2003), no. 1, 63–121. MR 2004457
- [FZ07] ———, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), no. 1, 112–164. MR 2295199
- [Kas01] R. M. Kashaev, *On the spectrum of Dehn twists in quantum Teichmüller theory*, Physics and combinatorics, 2000 (Nagoya), World Sci. Publ., River Edge, NJ, 2001, pp. 63–81. MR 1872252
- [Le16] Ian Le, *Higher laminations and affine buildings*, Geom. Topol. **20** (2016), no. 3, 1673–1735. MR 3523066
- [Pen87] R. C. Penner, *The decorated Teichmüller space of punctured surfaces*, Comm. Math. Phys. **113** (1987), no. 2, 299–339. MR 919235
- [Pen12] Robert C. Penner, *Decorated Teichmüller theory*, QGM Master Class Series, European Mathematical Society (EMS), Zürich, 2012, With a foreword by Yuri I. Manin. MR 3052157
- [PH92] R. C. Penner and J. L. Harer, *Combinatorics of train tracks*, Annals of Mathematics Studies, vol. 125, Princeton University Press, Princeton, NJ, 1992. MR 1144770
- [PP93] A. Papadopoulos and R. C. Penner, *The Weil-Petersson symplectic structure at Thurston’s boundary*, Trans. Amer. Math. Soc. **335** (1993), no. 2, 891–904. MR 1089420
- [Thu88] William P. Thurston, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), no. 2, 417–431. MR 956596